

On Sobolev extension domains in $\mathbf{R}^n$

By Pavel Shvartsman

Abstract

We describe a class of Sobolev W_p^k -extension domains $\Omega \subset \mathbf{R}^n$ determined by a certain inner subhyperbolic metric in Ω . This enables us to characterize finitely connected Sobolev W_p^1 -extension domains in $\mathbf{R}^2$ for each $p > 2$.

1. Introduction.

Let Ω be a domain in $\mathbf{R}^n$. This paper is devoted to the problem of extendability of functions from the Sobolev space $W_p^k(\Omega)$ to functions from $W_p^k(\mathbf{R}^n)$. We recall that, given $k \in \mathbf{N}$ and $p \in [1, \infty]$, the Sobolev space $W_p^k(\Omega)$, see e.g. Maz'ja [M], consists of all functions $f \in L_{1,loc}(\Omega)$ whose distributional partial derivatives on Ω of all orders up to k belong to $L_p(\Omega)$. $W_p^k(\Omega)$ is normed by

$$\|f\|_{W_p^k(\Omega)} := \sum \{\|D^\alpha f\|_{L_p(\Omega)} : |\alpha| \leq k\}.$$

A domain Ω in $\mathbf{R}^n$ is said to be a *Sobolev W_p^k -extension domain* if the following isomorphism

$$W_p^k(\Omega) = W_p^k(\mathbf{R}^n)|_\Omega$$

holds. In other words, Ω is a Sobolev extension domain (for the space $W_p^k(\mathbf{R}^n)$) if every Sobolev function $f \in W_p^k(\Omega)$ can be extended to a Sobolev W_p^k -function F defined on all of $\mathbf{R}^n$. For instance, Lipschitz domains (Calderón [C2], $1 < p < \infty$, Stein [St], $p = 1, \infty$) in $\mathbf{R}^n$ are W_p^k -extension domains for every $p \in [1, \infty]$ and every $k \in \mathbf{N}$. Jones [Jn] introduced a wider class of (ε, δ) -domains and proved that every (ε, δ) -domain is a Sobolev W_p^k -extension domain in $\mathbf{R}^n$ for every $k \geq 1$ and every $p \geq 1$. Burago and Maz'ya [BM], [M], Ch. 6, described extension domains for the space $BV(\mathbf{R}^n)$ of functions whose distributional derivatives of the first order are finite Radon measures.

Our main result is the following

Theorem 1.1 *Let $n < p < \infty$ and let Ω be a domain in $\mathbf{R}^n$. Suppose that there exist constants $C, \theta > 0$ such that the following condition is satisfied: for every $x, y \in \Omega$ such that $\|x - y\| \leq \theta$, there exists a rectifiable curve $\gamma \subset \Omega$ joining x to y such that*

$$\int_\gamma \text{dist}(z, \partial\Omega)^{\frac{1-n}{p-1}} ds(z) \leq C\|x - y\|^{\frac{p-n}{p-1}}. \quad (1.1)$$

Here $\partial\Omega$ denotes the boundary of Ω and ds denotes arc length measure.

Then Ω is a Sobolev W_q^k -extension domain for every $k \geq 1$ and every $q > p^*$ where $p^* \in (n, p)$ is a constant depending only on n, p and C .

Math Subject Classification 46E35

Key Words and Phrases Sobolev spaces, extension domains, subhyperbolic metric.

For $k = 1$ and $q > p$ this result has been proved by Koskela [K].

Observe that this theorem is also known for the case $p = \infty$ (with $p^* = q = \infty$). In that case every domain Ω satisfying inequality (1.1) is quasi-Euclidean, i.e., its inner metric is (locally) equivalent to the Euclidean distance. This case was studied by Whitney [W3] who proved that every quasi-Euclidean domain is a W_∞^k -extension domain for every $k \geq 1$.

Our next result, Theorem 1.2, relates to description of Sobolev extension domains in $\mathbf{R}^2$. The first result in this direction was obtained by Gol'dstein, Latfullin and Vodop'janov [GLV, GV1, GV2] who proved that a finitely connected bounded planar domain Ω is a Sobolev W_2^1 -extension domain if and only if Ω is an (ε, δ) -domain in $\mathbf{R}^2$ for some $\varepsilon, \delta > 0$. Maz'ja [M, MP] gave an example of a simply connected domain $\Omega \subset \mathbf{R}^2$ such that Ω is a W_p^1 -extension domain for every $p \in [1, 2)$, while $\mathbf{R}^2 \setminus \bar{\Omega}$ is a W_p^1 -extension domain for all $p > 2$. However Ω is not an (ε, δ) -domain for any ε and δ .

Buckley and Koskela [BKos] showed that if a finitely connected bounded domain $\Omega \subset \mathbf{R}^2$ is a Sobolev W_p^1 -extension domain for some $p > 2$, then there exists a constant $C > 0$ such that for every $x, y \in \Omega$ there exists a rectifiable curve $\gamma \subset \Omega$ satisfying inequality (1.1) (with $n = 2$). Combining this result with Theorem 1.1, we obtain the following

Theorem 1.2 *Let $2 < p < \infty$ and let Ω be a finitely connected bounded planar domain. Then Ω is a Sobolev W_p^1 -extension domain if and only if for some $C > 0$ the following condition is satisfied: for every $x, y \in \Omega$ there exists a rectifiable curve $\gamma \subset \Omega$ joining x to y such that*

$$\int_\gamma \text{dist}(z, \partial\Omega)^{\frac{1}{1-p}} ds(z) \leq C \|x - y\|^{\frac{p-2}{p-1}}.$$

We note that this result is also true for the case $p = \infty$ and then the space W_∞^1 can even be replaced by W_∞^k for arbitrary $k \geq 1$. This follows from the aforementioned theorem of Whitney [W3] combined with a result of Zobin [Zob2] which states that every finitely connected bounded planar W_∞^k -extension domain is quasi-Euclidean. Zobin [Zob1] also showed that for every $k \geq 2$ there exists a bounded planar W_∞^k -extension domain which is not quasi-Euclidean.

Let us briefly indicate the main ideas of our approach for the case $k = 1$, i.e., for the Sobolev space $W_p^1(\mathbf{R}^n)$. Recall that, when $p > n$, it follows from the Sobolev embedding theorem that every function $f \in W_p^1(\Omega)$, $p > n$, can be redefined, if necessary, on a subset of Ω of Lebesgue measure zero so that it satisfies a *local* Hölder condition of order $\alpha := 1 - \frac{n}{p}$ on Ω : i.e., for every ball $B \subset \Omega$

$$|f(x) - f(y)| \leq C(n, p) \|f\|_{W_p^1(\Omega)} \|x - y\|^{1-\frac{n}{p}}, \quad x, y \in B. \quad (1.2)$$

We will identify each element of $W_p^1(\Omega)$ with its unique continuous representative. Thus we will be able to restrict our attention to the case of continuous Sobolev functions.

Following Buckley and Stanoyevitch [BSt3], given $\alpha \in [0, 1]$ and a rectifiable curve $\gamma \subset \Omega$, we define the *subhyperbolic length* of γ by

$$\text{len}_{\alpha, \Omega}(\gamma) := \int_\gamma \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z).$$

Then we let $d_{\alpha,\Omega}$ denote the corresponding *subhyperbolic metric* on Ω given, for each $x, y \in \Omega$, by

$$d_{\alpha,\Omega}(x, y) := \inf_{\gamma} \text{len}_{\alpha,\Omega}(\gamma) \quad (1.3)$$

where the infimum is taken over all rectifiable curves $\gamma \subset \Omega$ joining x to y .

The metric $d_{\alpha,\Omega}$ was introduced and studied by Gehring and Martio in [GM]. See also [AHHL, L, BKos] for various further results using this metric. Note also that $\text{len}_{0,\Omega}$ and $d_{0,\Omega}$ are the well-known *quasihyperbolic length* and *quasihyperbolic distance*, and $d_{1,\Omega}$ is the *inner (or geodesic) metric* on Ω .

The subhyperbolic metric $d_{\alpha,\Omega}$ with $\alpha = (p - n)/(p - 1)$ arises naturally in the study of Sobolev $W_p^1(\Omega)$ -functions for $p > n$. In particular, Buckley and Stanoyevitch [BSt2] proved that the local Hölder condition (1.2) is equivalent to the following Hölder-type condition

$$|f(x) - f(y)| \leq C(n, p) \|f\|_{W_p^1(\Omega)} \{d_{\alpha,\Omega}(x, y)^{1-\frac{1}{p}} + \|x - y\|^{1-\frac{n}{p}}\}, \quad x, y \in \Omega, \quad (1.4)$$

with $\alpha = (p - n)/(p - 1)$.

In turn, since any extension $F \in W_p^1(\mathbf{R}^n)$ of f satisfies the global Hölder condition

$$|F(x) - F(y)| \leq C(n, p) \|F\|_{W_p^1(\mathbf{R}^n)} \|x - y\|^{1-\frac{n}{p}}, \quad x, y \in \mathbf{R}^n,$$

we have

$$|f(x) - f(y)| \leq C(n, p) \|F\|_{W_p^1(\mathbf{R}^n)} \|x - y\|^{1-\frac{n}{p}}, \quad x, y \in \Omega. \quad (1.5)$$

Of course the conditions (1.4) and (1.5) with $\|f\|_{W_p^1(\Omega)}$ and $\|F\|_{W_p^1(\mathbf{R}^n)}$ replaced by unspecified constants are not equivalent to membership of f in $W_p^1(\Omega)$ or in $W_p^1(\mathbf{R}^n)|_{\Omega}$ respectively. However the preceding remarks suggest that a reasonable property which might perhaps be necessary or perhaps sufficient for a domain Ω to be a Sobolev extension domain could be this: Whenever a function $f : \Omega \rightarrow \mathbf{R}$ satisfies

$$|f(x) - f(y)| \leq d_{\alpha,\Omega}(x, y)^{1-\frac{1}{p}} + \|x - y\|^{1-\frac{n}{p}}$$

for all $x, y \in \Omega$ and $\alpha = (p - n)/(p - 1)$ then it also satisfies

$$|f(x) - f(y)| \leq C(n, p) \|x - y\|^{1-\frac{n}{p}}$$

for all $x, y \in \Omega$ and for some constant $C(n, p)$ depending only on n and p .

One would like to have a simpler condition on Ω which would be sufficient to imply the above “reasonable property”. It is clear that the following property, which has already been considered and studied by other authors, namely

$$d_{\alpha,\Omega}(x, y)^{1-\frac{1}{p}} \leq C \|x - y\|^{1-\frac{n}{p}} \quad \text{for all } x, y \in \Omega \text{ and } \alpha = (p - n)/(p - 1)$$

or, equivalently, $d_{\alpha,\Omega}(x, y) \leq C \|x - y\|^\alpha$ for all $x, y \in \Omega$, is such a condition.

These considerations lead us to work with a certain class of domains, essentially those which were introduced in [GM]. In our context here, it seems convenient to use terminology different from that of [GM] and other papers.

Definition 1.3 For each $\alpha \in (0, 1]$, the domain $\Omega \subset \mathbf{R}^n$ is said to be α -subhyperbolic if there exist constants $C_{\alpha, \Omega} > 0$ and $\theta_{\alpha, \Omega} > 0$ such that

$$d_{\alpha, \Omega}(x, y) \leq C_{\alpha, \Omega} \|x - y\|^\alpha$$

for every $x, y \in \Omega$ satisfying $\|x - y\| \leq \theta_{\alpha, \Omega}$.

We denote the class of α -subhyperbolic domains in $\mathbf{R}^n$ by $U_\alpha(\mathbf{R}^n)$.

In [GM] and also in [L] these domains are called “ Lip_α -extension domains”. (This name is derived from the fact that $\Omega \in U_\alpha(\mathbf{R}^n)$ iff all functions which are *locally* Lipschitz of order α on Ω are Lipschitz of order α on Ω .) These domains have also been studied in [BSt2, BSt, BSt3] where they are called “ $\alpha - m$ cigar domains”, and in [BKos] where they are termed “local weak α -cigar domains”.

Now Theorem 1.1 can be reformulated as follows: For each $p > n$ and for each $\frac{p-n}{p-1}$ -subhyperbolic domain Ω in $\mathbf{R}^n$, there exists a constant $p^* \in (n, p)$ depending only on n , p and Ω , such that Ω is a Sobolev W_q^k -extension domain for every $q \geq p^*$.

In turn, Theorem 1.2 admits the following reformulation: For each $p > 2$, a finitely connected bounded domain $\Omega \subset \mathbf{R}^2$ is a Sobolev W_p^1 -extension domain if and only if Ω is a $\frac{p-2}{p-1}$ -subhyperbolic domain.

The family $\{U_\alpha(\mathbf{R}^n) : \alpha \in (0, 1]\}$ is an “increasing family”, i.e.,

$$U_{\alpha'}(\mathbf{R}^n) \subset U_{\alpha''}(\mathbf{R}^n) \quad \text{whenever } 0 < \alpha' < \alpha'' \leq 1,$$

see, e.g. [BKos]. Lappalainen [L] proved that

$$U_\alpha(\mathbf{R}^n) \subsetneq \bigcap_{\alpha < \tau \leq 1} U_\tau(\mathbf{R}^n) \quad \text{for every } \alpha \in (0, 1).$$

This last result motivates our discussion presented in Section 2, which is devoted to the following question: Does the equality

$$U_\alpha(\mathbf{R}^n) = \bigcup_{0 < \tau < \alpha} U_\tau(\mathbf{R}^n) \tag{1.6}$$

hold? In other words, do α -subhyperbolic domains have the following “self-improving” property that whenever Ω is an α -subhyperbolic domain in $\mathbf{R}^n$ for some $\alpha \in (0, 1)$, it is also τ -subhyperbolic for some positive τ which is *strictly less* than α ? (Of course, τ can depend on Ω).

We do not know the answer to this question in general. We do know that the answer is affirmative for an arbitrary finitely connected bounded domain $\Omega \in U_\alpha(\mathbf{R}^2)$, $\alpha \in (0, 1)$, as it follows from Theorem 1.1 and Theorem 1.2. We also know that for a certain subfamily of $U_\alpha(\mathbf{R}^n)$, the so-called *strongly* α -subhyperbolic domains (Definition 2.4) the answer to the above question is affirmative. (See Proposition 2.6.) It should be pointed out that we have no examples of subhyperbolic domains which are not strongly subhyperbolic.

We are able to show that the following weaker version of the self-improving property (1.6) holds for an *arbitrary* subhyperbolic domain in $\mathbf{R}^n$.

Theorem 1.4 *Let $\alpha \in (0, 1)$ and let Ω be an α -subhyperbolic domain in $\mathbf{R}^n$. There exist a constant $\alpha^*, 0 < \alpha^* < \alpha$, and constants $\theta, C > 0$ such that the following is true:*

For every $\varepsilon > 0$ and every $x, y \in \Omega$, $\|x - y\| \leq \theta$, there exist a rectifiable curve $\Gamma \subset \Omega$ joining x to y and a subset $\tilde{\Gamma} \subset \Gamma$ consisting of a finite number of arcs such that the following conditions are satisfied:

(i). For every $\tau \in [\alpha^*, \alpha]$

$$\int_{\tilde{\Gamma}} \text{dist}(z, \partial\Omega)^{\tau-1} ds(z) \leq C\|x - y\|^\tau. \quad (1.7)$$

In addition, for every ball B centered in $\tilde{\Gamma}$ of radius at most $\|x - y\|$,

$$\text{diam } B \leq C \text{ length}(B \cap \tilde{\Gamma}). \quad (1.8)$$

(ii). We have $\text{length}(\Gamma) \leq C\|x - y\|$ and

$$\text{length}(\Gamma \setminus \tilde{\Gamma}) < \varepsilon. \quad (1.9)$$

Moreover,

$$\int_{\Gamma \setminus \tilde{\Gamma}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C\|x - y\|^\alpha. \quad (1.10)$$

The constants α^*, θ and C depend only on n, α , and the constants $C_{\alpha, \Omega}$ and $\theta_{\alpha, \Omega}$ introduced in Definition 1.3.

The proof of this result, presented in Section 2, is based on the reverse Hölder inequality for m -dyadic A_1 -weights. (See Melas [Mel].)

Theorem 1.4 is an important ingredient in the proof of the extension Theorem 1.1. It enables us to prove the following version of the *Sobolev-Poincaré inequality* for subhyperbolic domains (for $p > n$ and $k \geq 1$): Let Ω be an α -subhyperbolic domain in $\mathbf{R}^n$ with $\alpha = (p - n)/(p - 1)$. Given $f \in C^{k-1}(\Omega)$ and $x \in \Omega$ we let $T_x^{k-1}(f)$ denote the Taylor polynomial of f at x of degree at most $k - 1$. We prove that there exists $p^* \in (n, p)$ and constants $\theta, \lambda, C > 0$ such that for every function $f \in C^{k-1}(\Omega) \cap W_p^k(\Omega)$ and every $x, y \in \Omega$, $\|x - y\| \leq \theta$, the following inequality

$$|f(y) - T_x^{k-1}(f)(y)| \leq C\|x - y\|^{k - \frac{n}{p^*}} \left(\int_{B \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}$$

holds. Here $B = B(x, \lambda\|x - y\|)$ is the ball centered at x of radius $r = \lambda\|x - y\|$. This inequality is a particular case of Theorem 3.1 which we prove in Section 3.

In Section 4 we prove a corollary of this result related to the sharp maximal function

$$f_{k, \Omega}^\#(x) := \sup_{r > 0} r^{-k} \inf_{P \in \mathcal{P}_{k-1}} \frac{1}{|B(x, r)|} \int_{B(x, r) \cap \Omega} |f - P| dx, \quad x \in \Omega.$$

Here $\mathcal{P}_{k-1}$ is the space of polynomials of degree at most $k-1$ defined on $\mathbf{R}^n$ and $|B(x, r)|$ is the Lebesgue measure of the ball $B(x, r)$. We show that for every $f \in W_p^k(\Omega)$ and every $x \in \Omega$ the following inequality

$$f_{k,\Omega}^\sharp(x) \leq C \left\{ (\mathcal{M}[(\|\nabla^k f\|^\wedge)^{p^*}](x))^{\frac{1}{p^*}} + \mathcal{M}[f^\wedge](x) \right\} \quad (1.11)$$

holds. Here $\mathcal{M}$ denotes the Hardy-Littlewood maximal function and the symbol $g^\wedge$ stands for the extension *by zero* of a function from Ω to all of $\mathbf{R}^n$.

The sharp maximal function is a useful tool in the study of Sobolev functions. In [C] Calderón proved that, for $p > 1$, a function f is in $W_p^k(\mathbf{R}^n)$ if and only if f and $f_{k,\mathbf{R}^n}^\sharp$ are both in $L_p(\mathbf{R}^n)$. In [S4] this description has been generalized to the case of the so-called *regular* subsets of $\mathbf{R}^n$, i.e., the sets S such that $|B \cap S| \sim |B|$ for all balls B centered in S of radius at most 1. We proved in [S4] that if S is regular and $f \in L_p(S)$, $p > 1$, then f can be extended to a function $F \in W_p^k(\mathbf{R}^n)$ if and only if its sharp maximal function $f_{k,S}^\sharp \in L_p(S)$. (For the case $k = 1$ see also [S3, HKT, HKT1]. Observe that every Sobolev W_p^1 -extension domain is a regular subset of $\mathbf{R}^n$, see Hajlasz, Koskela and Tuominen [HKT]. In [S7] we present a description of the trace space $W_p^1(\mathbf{R}^n)|_S$, $p > n$, for an *arbitrary* set $S \subset \mathbf{R}^n$ via an L_∞ -version of the sharp maximal function).

Every subhyperbolic domain is a regular set, as shown in Lemma 2.3. So, in order to prove, for some given $q > p^*$, that a function $f \in W_q^k(\Omega)$ extends to a Sobolev W_q^k -function on $\mathbf{R}^n$, it suffices to show that $f_{k,\Omega}^\sharp \in L_q(\Omega)$. We do this by applying the Hardy-Littlewood maximal theorem to inequality (1.11). This gives us the inequality $\|f_{k,\Omega}^\sharp\|_{L_q(\Omega)} \leq C\|f\|_{W_q^k(\Omega)}$ which completes the proof of Theorem 1.1.

Acknowledgement. The author is greatly indebted to Michael Cwikel, Charles Fefferman, Vladimir Maz'ya and Naum Zobin for interesting discussions and helpful suggestions and remarks.

2. Subhyperbolic domains: intrinsic metrics and self-improvement.

Throughout the paper $C, C_1, C_2, \dots$ will be generic positive constants which depend only on parameters determining sets (say, n, α , the constants $C_{\alpha,\Omega}$ or $\theta_{\alpha,\Omega}$, etc.) or function spaces (p, q , etc). These constants can change even in a single string of estimates. The dependence of a constant on certain parameters is expressed, for example, by the notation $C = C(n, p)$.

The Lebesgue measure of a measurable set $A \subset \mathbf{R}^n$ will be denoted by $|A|$. Given subsets $A, B \subset \mathbf{R}^n$, we put $\text{diam } A := \sup\{\|a - a'\| : a, a' \in A\}$ and

$$\text{dist}(A, B) := \inf\{\|a - b\| : a \in A, b \in B\}.$$

For $x \in \mathbf{R}^n$ we also set $\text{dist}(x, A) := \text{dist}(\{x\}, A)$.

Let $\gamma : [a, b] \rightarrow \mathbf{R}^n$ be a curve in $\mathbf{R}^n$, and let $u = \gamma(t_1), v = \gamma(t_2)$ where $a \leq t_1 < t_2 \leq b$. By γ_{uv} we denote the arc of γ joining u to v .

We will be needed the following auxiliary lemma.

Lemma 2.1 (i). Let $x, y \in \Omega$ and let

$$\max(\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)) \leq 2\|x - y\|. \quad (2.1)$$

Let γ be a rectifiable curve joining x to y in Ω . Assume that for some $\alpha \in (0, 1)$ and $C > 0$ the following inequality

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C\|x - y\|^{\alpha} \quad (2.2)$$

holds. Then

$$\text{length}(\gamma) \leq 2e^C\|x - y\|.$$

(ii). Let $x, y \in \Omega$ and let

$$\max(\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)) > 2\|x - y\|. \quad (2.3)$$

Then the line segment $[x, y] \subset \Omega$ and for every $\beta \in (0, 1]$ we have

$$\int_{[x, y]} \text{dist}(z, \partial\Omega)^{\beta-1} ds(z) \leq \|x - y\|^{\beta}. \quad (2.4)$$

Proof. (i). Let us parameterize γ by arclength; thus we identify γ with a function $\gamma : [0, \ell] \rightarrow \Omega$ satisfying $\gamma(0) = x, \gamma(\ell) = y$. Now (2.2) is equivalent to

$$\int_0^{\ell} \text{dist}(\gamma(t), \partial\Omega)^{\alpha-1} dt \leq C\|x - y\|^{\alpha}.$$

Since $\text{dist}(\cdot, \partial\Omega)$ is a Lipschitz function on $\mathbf{R}^n$,

$$\text{dist}(u, \partial\Omega) \leq \text{dist}(v, \partial\Omega) + \|u - v\|, \quad u, v \in \Omega, \quad (2.5)$$

so that for every $t \in (0, \ell]$

$$\text{dist}(\gamma(t), \partial\Omega) \leq \text{dist}(x, \partial\Omega) + \|x - \gamma(t)\|.$$

Since γ is parameterized by arclength,

$$\|x - \gamma(t)\| \leq \text{length}(\gamma_{x\gamma(t)}) = t,$$

so that

$$\text{dist}(\gamma(t), \partial\Omega) \leq \text{dist}(x, \partial\Omega) + t, \quad t \in [0, \ell].$$

This inequality and (2.2) imply

$$\begin{aligned} C\|x - y\|^{\alpha} &\geq \int_0^{\ell} \text{dist}(\gamma(t), \partial\Omega)^{\alpha-1} dt \geq \int_0^{\ell} (\text{dist}(x, \partial\Omega) + t)^{\alpha-1} dt \\ &= \alpha^{-1}((\text{dist}(x, \partial\Omega) + \ell)^{\alpha} - \text{dist}(x, \partial\Omega)^{\alpha}) \\ &\geq \alpha^{-1}(\ell^{\alpha} - \text{dist}(x, \partial\Omega)^{\alpha}). \end{aligned}$$

But $2\|x - y\| > \text{dist}(x, \partial\Omega)$ so that

$$C\|x - y\|^\alpha \geq \alpha^{-1}(\ell^\alpha - (2\|x - y\|)^\alpha).$$

Hence

$$\ell \leq (\alpha C + 2^\alpha)^{\frac{1}{\alpha}}\|x - y\| \leq 2e^C\|x - y\|$$

proving (i).

(ii). Clearly, (2.3) implies $[x, y] \subset \Omega$. Prove (2.4).

We may assume that $\text{dist}(x, \partial\Omega) > 2\|x - y\|$. Also note that $\|x - z\| \leq \|x - y\|$ for every $z \in [x, y]$. These inequalities and (2.5) imply the following:

$$\frac{1}{2} \text{dist}(x, \partial\Omega) \leq \text{dist}(x, \partial\Omega) - \|x - y\| \leq \text{dist}(x, \partial\Omega) - \|x - z\| \leq \text{dist}(z, \partial\Omega).$$

Hence,

$$\begin{aligned} \int_{[x,y]} \text{dist}(z, \partial\Omega)^{\beta-1} ds(z) &\leq \int_{[x,y]} 2^{1-\beta} \text{dist}(x, \partial\Omega)^{\beta-1} ds(z) \\ &= 2^{1-\beta} \|x - y\| \text{dist}(x, \partial\Omega)^{\beta-1} \\ &\leq 2^{1-\beta} \|x - y\| (2\|x - y\|)^{\beta-1} = \|x - y\|^\beta \end{aligned}$$

proving the lemma. $\square$

Lemma 2.2 *Let $x, y \in \Omega$ and let $\gamma \subset \Omega$ be a rectifiable curve joining x to y . Suppose that for some $\alpha \in (0, 1)$ and $C \geq 1$ the following inequality*

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C \text{length}^\alpha(\gamma) \quad (2.6)$$

holds. Then

(i). *There exists a point $\bar{z} \in \gamma$ such that*

$$\text{length}(\gamma) \leq C^{\frac{1}{1-\alpha}} \text{dist}(\bar{z}, \partial\Omega)$$

(ii). *We have*

$$\frac{1}{\text{length}(\gamma)} \int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq 2C \inf_{z \in \gamma} \text{dist}(z, \partial\Omega)^{\alpha-1}.$$

Proof. (i). Put $\ell := \text{length}(\gamma)$. Let $\bar{z}$ be a point in γ such that

$$\max\{\text{dist}(z, \partial\Omega) : z \in \gamma\} = \text{dist}(\bar{z}, \partial\Omega).$$

Then

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \geq \int_{\gamma} \text{dist}(\bar{z}, \partial\Omega)^{\alpha-1} ds(z) = \ell \text{dist}(\bar{z}, \partial\Omega)^{\alpha-1}$$

so that, by (2.6),

$$\ell \operatorname{dist}(z, \partial\Omega)^{\alpha-1} \leq C\ell^\alpha.$$

Hence

$$\ell \leq C^{\frac{1}{1-\alpha}} \operatorname{dist}(z, \partial\Omega)$$

proving (i).

(ii). Put $w(z) := \operatorname{dist}(z, \partial\Omega)$. Then, by (2.6),

$$\frac{1}{\ell} \int_{\gamma} w(z)^{\alpha-1} ds(z) \leq \ell^{-1} (C \ell^\alpha) = C \ell^{\alpha-1}. \quad (2.7)$$

For every $z_1, z_2 \in \gamma$ we have

$$|w(z_1) - w(z_2)| = |\operatorname{dist}(z_1, \partial\Omega) - \operatorname{dist}(z_2, \partial\Omega)| \leq \|z_1 - z_2\| \leq \ell$$

so that

$$\max_{z \in \gamma} w(z) \leq \min_{z \in \gamma} w(z) + \ell. \quad (2.8)$$

Let us consider two cases. First suppose that $\max_{z \in \gamma} w(z) \leq 2\ell$. Since $\alpha \in (0, 1)$, we obtain

$$\ell^{\alpha-1} \leq 2^{1-\alpha} \min_{z \in \gamma} w(z)^{\alpha-1}$$

so that, by (2.7),

$$\frac{1}{\ell} \int_{\gamma} w(z)^{\alpha-1} ds(z) \leq 2^{1-\alpha} C \min_{z \in \gamma} w(z)^{\alpha-1} \leq 2C \min_{z \in \gamma} w(z)^{\alpha-1}.$$

Now assume that $2\ell < \max_{z \in \gamma} w(z)$. Then, by (2.8),

$$\max_{z \in \gamma} w(z) \leq \min_{z \in \gamma} w(z) + \frac{1}{2} \max_{z \in \gamma} w(z)$$

so that $\max_{z \in \gamma} w(z) \leq 2 \min_{z \in \gamma} w(z)$. Hence

$$\max_{z \in \gamma} w(z)^{\alpha-1} \leq 2^{1-\alpha} \min_{z \in \gamma} w(z)^{\alpha-1} \leq 2 \min_{z \in \gamma} w(z)^{\alpha-1}.$$

Finally, we have

$$\frac{1}{\ell} \int_{\gamma} w(z)^{\alpha-1} ds(z) \leq \max_{z \in \gamma} w(z)^{\alpha-1} \leq 2 \min_{z \in \gamma} w(z)^{\alpha-1}.$$

The lemma is proved.

This lemma implies the following important property of subhyperbolic domains.

Lemma 2.3 *Let $\alpha \in (0, 1)$ and let Ω be an α -subhyperbolic domain.*

There exist constant $\delta > 0$ and $\sigma \in (0, 1]$ depending only on $n, \alpha, C_{\alpha, \Omega}$ and $\theta_{\alpha, \Omega}$ such that every ball B centered in Ω of diameter at most δ contains a ball $B' \subset \Omega$ of diameter at least $\sigma \operatorname{diam} B$.

Proof. Let $\delta := \min\{\theta_{\alpha,\Omega}, \frac{1}{2} \text{diam } \Omega\}$. Let $B = B(x, r)$ be a ball with center in $x \in \Omega$ and radius $r \leq \delta$. Put $\tilde{r} := r/(8e^{C_{\alpha,\Omega}})$. Since $\tilde{r} \leq \delta \leq \frac{1}{2} \text{diam } \Omega$, there exists a point $a \in \Omega$ such that $\|x - a\| > \tilde{r}$. Let $\Gamma \subset \Omega$ be a curve joining x to a . Since $a \notin B(x, r)$, we have $\Gamma \cap \partial(B(x, r)) \neq \emptyset$ so that there exists a point $b \in \Omega$ such that $\|x - b\| = \tilde{r}$.

If

$$\max(\text{dist}(x, \partial\Omega), \text{dist}(b, \partial\Omega)) > 2\|x - b\| = 2\tilde{r},$$

then either $B(x, \tilde{r}) \subset B(x, r) \cap \Omega$ or $B(b, \tilde{r}) \subset B(x, r) \cap \Omega$, so that in this case the ball B' exists.

Suppose that

$$\max(\text{dist}(x, \partial\Omega), \text{dist}(b, \partial\Omega)) \leq 2\|x - b\| = 2\tilde{r},$$

Since $\tilde{r} \leq \delta \leq \theta_{\alpha,\Omega}$, there exists a curve $\gamma \subset \Omega$ joining x to b such that

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C_{\alpha,\Omega} \|x - b\|^{\alpha}.$$

(We may assume that $C_{\alpha,\Omega} \geq 1$.) By Lemma 2.1, part (i),

$$\text{length}(\gamma) \leq 2e^{C_{\alpha,\Omega}} \|x - b\| = 2e^{C_{\alpha,\Omega}} \tilde{r} = r/4.$$

Moreover, by part (ii) of Lemma 2.2, there exists a point $\bar{z} \in \gamma$ such that

$$\text{length}(\gamma) \leq C_{\alpha,\Omega}^{\frac{1}{1-\alpha}} \text{dist}(\bar{z}, \partial\Omega).$$

Hence,

$$\tilde{r} = \|x - b\| \leq \text{length}(\gamma) \leq C_{\alpha,\Omega}^{\frac{1}{1-\alpha}} \text{dist}(\bar{z}, \partial\Omega).$$

Put $r' := \tilde{r}/(2C_{\alpha,\Omega}^{\frac{1}{1-\alpha}})$ and $B' := B(z, r')$. Then, $r' \leq \tilde{r} \leq r/4$ (recall that $C_{\alpha,\Omega} \geq 1$) and $r' < \text{dist}(\bar{z}, \partial\Omega)$.

Hence, $B(z, \tilde{r}) \subset \Omega$. On the other hand, $\|x - z\| \leq \text{length}(\gamma) \leq r/4$ so that $B' \subset B(x, r/2) \subset B$. The lemma is proved. $\square$

Before to present the proof of Theorem 1.4 let us demonstrate its main ideas for a family of the so-called strongly subhyperbolic domains in $\mathbf{R}^n$

Definition 2.4 Let $\alpha \in (0, 1]$. A domain $\Omega \subset \mathbf{R}^n$ is said to be *strongly α -subhyperbolic* if there exist constants $C, \theta > 0$ such that every $x, y \in \Omega$, $\|x - y\| \leq \theta$, can be joined by a rectifiable curve $\gamma \subset \Omega$ satisfying the following condition: for every $u, v \in \gamma$

$$\int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C \|u - v\|^{\alpha}. \quad (2.9)$$

Remark 2.5 Given $x, y \in \Omega$ a rectifiable curve $\gamma \subset \Omega$ joining x to y is said to be $d_{\alpha,\Omega}$ -geodesic if

$$d_{\alpha,\Omega}(x, y) = \text{len}_{\alpha,\Omega}(\gamma) := \int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z).$$

(See definition (1.3).)

Clearly, if Ω is α -subhyperbolic and for every $x, y \in \Omega$ there exists $d_{\alpha, \Omega}$ -geodesic, then Ω is strongly α -subhyperbolic. In fact, in this case every arc of $d_{\alpha, \Omega}$ -geodesic curve is $d_{\alpha, \Omega}$ -geodesic as well so that inequality (2.9) holds.

However, for every $\alpha \in (0, 1]$ there exists a domain $\Omega \in \mathbf{R}^n$ and $x, y \in \Omega$ such that $d_{\alpha, \Omega}$ -geodesic for x, y does not exist. This is trivial for $\alpha = 1$, i.e., for quasi-Euclidean domains. For the case $\alpha \in (0, 1)$ see [BSt2].

Let us slightly generalize this example. Fix $C \geq 1$. We say that a rectifiable curve $\gamma \subset \Omega$ joining x to y is $(C, d_{\alpha, \Omega})$ -geodesic if for every $u, v \in \gamma$ the following inequality

$$\text{len}_{\alpha, \Omega}(\gamma_{uv}) \leq C d_{\alpha, \Omega}(u, v).$$

holds. Clearly, a rectifiable curve γ is $(1, d_{\alpha, \Omega})$ -geodesic iff it is $d_{\alpha, \Omega}$ -geodesic. Moreover, if $\Omega \in U_{\alpha}(\mathbf{R}^n)$ and for every $x, y \in \Omega$, $\|x - y\| \leq \theta$, there exists $(C, d_{\alpha, \Omega})$ -geodesic joining x to y in Ω , then Ω is strongly α -subhyperbolic.

This observation motivates the following question: Let Ω be a domain in $\mathbf{R}^n$ and let $\alpha \in (0, 1]$. Does there exist a constant $C = C_{\Omega} > 1$ such that every two points $x, y \in \Omega$ can be joined by a $(C, d_{\alpha, \Omega})$ -geodesic curve? Even for the quasi-Euclidean domains, i.e., for $\alpha = 1$, we do not know the answer to this question. $\square$

Proposition 2.6 *Let $\alpha \in (0, 1)$ and let Ω be a strongly α -subhyperbolic domain in $\mathbf{R}^n$. Then Ω is τ -subhyperbolic for some $\tau \in (0, \alpha)$.*

Proof. Since Ω is strongly α -subhyperbolic, there exist constants $\theta > 0$ and $C \geq 1$ such that every $x, y \in \Omega$, $\|x - y\| \leq \theta$, can be joined by a rectifiable curve $\gamma \subset \Omega$ satisfying the following condition: for every $u, v \in \gamma$

$$\int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C \|u - v\|^{\alpha}.$$

In particular,

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C \|x - y\|^{\alpha}. \quad (2.10)$$

Let $\ell := \text{length}(\gamma)$. We parameterize γ by arclength: thus $\gamma : [0, \ell] \rightarrow \Omega$, $\gamma(0) = x$, $\gamma(\ell) = y$. Let $u = \gamma(t_1)$, $v = \gamma(t_2)$ where $0 < t_1 < t_2 \leq \ell$ (recall that by γ_{uv} we denote the arc of γ joining u to v).

Applying part (ii) of Lemma 2.2 to the arc γ_{uv} , we obtain

$$\frac{1}{\text{length}(\gamma_{uv})} \int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq 2C \inf_{z \in \gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1},$$

or, in the parametric form,

$$\frac{1}{t_2 - t_1} \int_{t_1}^{t_2} \text{dist}(\gamma(t), \partial\Omega)^{\alpha-1} dt \leq 2C \inf_{t \in [t_1, t_2]} \text{dist}(\gamma(t), \partial\Omega)^{\alpha-1}. \quad (2.11)$$

Put

$$w(t) := \text{dist}(\gamma(t), \partial\Omega).$$

By (2.11), the function $w = w(t)$ has the following property: for every subinterval $I \subset [0, \ell]$

$$\frac{1}{|I|} \int_I w(t)^{\alpha-1} dt \leq 2C \inf_I w(t)^{\alpha-1}.$$

Thus the function $h := w^{\alpha-1}$ is a Muckenhoupt's $\mathcal{A}_1$ -weight on $[0, \ell]$, see, e.g., [GR], so that h satisfies the reverse Hölder inequality on $[0, \ell]$ (see Mackenhaupt [Mac], Gehring [G], Coiffman and Fefferman [CF]): There exist constants $\tilde{q} > 1$ and $C_1 \geq 1$ (depending only on C) such that

$$\left(\frac{1}{\ell} \int_0^\ell h^{\tilde{q}}(t) dt \right)^{1/\tilde{q}} \leq C_1 \frac{1}{\ell} \int_0^\ell h(t) dt.$$

By (2.10),

$$\int_0^\ell w^{\alpha-1}(t) dt \leq C \|x - y\|^\alpha$$

so that

$$\left(\frac{1}{\ell} \int_0^\ell w^{(\alpha-1)\tilde{q}}(t) dt \right)^{1/\tilde{q}} \leq C_1 \frac{1}{\ell} \int_0^\ell w^{\alpha-1}(t) dt \leq C_1 C \frac{1}{\ell} \|x - y\|^\alpha.$$

We put $q := \min\{\tilde{q}, \frac{1-\alpha/2}{1-\alpha}\}$ and $\tau = q(\alpha - 1) + 1$. Clearly, $1 < q \leq \tilde{q}$ and $0 < \tau < \alpha$. Hence,

$$\begin{aligned} \left(\frac{1}{\ell} \int_0^\ell w^{\tau-1}(t) dt \right)^{1/q} &= \left(\frac{1}{\ell} \int_0^\ell w^{q(\alpha-1)}(t) dt \right)^{1/q} \\ &\leq \left(\frac{1}{\ell} \int_0^\ell w^{\tilde{q}(\alpha-1)}(t) dt \right)^{1/\tilde{q}} \leq C_2 \frac{1}{\ell} \|x - y\|^\alpha \end{aligned}$$

where $C_2 := C_1 C$. Finally, we obtain

$$\begin{aligned} \int_\gamma \text{dist}(z, \partial\Omega)^{\tau-1} ds(z) &= \int_0^\ell w^{\tau-1}(t) dt \leq \frac{C_2^q}{\ell^{q-1}} \|x - y\|^{\alpha q} \\ &\leq C_2^q \|x - y\|^{\alpha q - q + 1} = C_2^q \|x - y\|^\tau \end{aligned}$$

proving that Ω is a τ -subhyperbolic domain. $\square$

Proof of Theorem 1.4. Let $\varepsilon > 0$ and let $\Omega \in U_\alpha(\mathbf{R}^n)$. We will assume that the constant $C_{\alpha, \Omega} \geq 1$. Put

$$\theta := \frac{1}{2} e^{-2C_{\alpha, \Omega}} \theta_{\alpha, \Omega}$$

and fix $x, y \in \Omega$ such that $\|x - y\| \leq \theta$.

By part (ii) of Lemma 2.1, if inequality (2.3) is satisfied, then the statement of Theorem 1.4 is true with $\Gamma = \tilde{\Gamma} = [x, y]$ and any $\alpha^* \in (0, \alpha)$.

Now suppose that x, y satisfy inequality (2.1), i.e.,

$$\max(\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)) \leq 2\|x - y\|.$$

Lemma 2.7 *Let $\Omega \in U_\alpha(\mathbf{R}^n)$ and let $x, y \in \Omega$, $\|x - y\| \leq \theta$. Let $0 < \delta \leq d_{\alpha, \Omega}(x, y)$ and let $\gamma \subset \Omega$ be a rectifiable curve joining x to y such that*

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) < d_{\alpha, \Omega}(x, y) + \delta. \quad (2.12)$$

Then:

(i). We have

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq 2C_{\alpha, \Omega} \|x - y\|^\alpha$$

and $\text{length}(\gamma) \leq 2e^{2C_{\alpha, \Omega}} \|x - y\|$.

(ii). For every $u, v \in \gamma$ such that

$$\text{length}(\gamma_{uv}) \geq \delta^{\frac{1}{\alpha}} \quad (2.13)$$

the following inequality

$$\int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq 2C_{\alpha, \Omega} \text{length}^\alpha(\gamma_{uv})$$

holds.

Proof. First prove that

$$\int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) < d_{\alpha, \Omega}(u, v) + \delta. \quad (2.14)$$

In fact, assume that

$$d_{\alpha, \Omega}(u, v) + \delta \leq \int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z).$$

Then

$$\begin{aligned} \int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) &= \int_{\gamma_{xu}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) + \int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \\ &\quad + \int_{\gamma_{vy}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \\ &\geq d_{\alpha, \Omega}(x, u) + (d_{\alpha, \Omega}(u, v) + \delta) + d_{\alpha, \Omega}(v, y) \end{aligned}$$

so that, by the triangle inequality for the metric $d_{\alpha, \Omega}$,

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \geq d_{\alpha, \Omega}(x, y) + \delta$$

which contradicts inequality (2.12).

Since $0 < \delta \leq d_{\alpha,\Omega}(x, y)$, by (2.12),

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) < 2d_{\alpha,\Omega}(x, y).$$

Since $\theta \leq \theta_{\alpha,\Omega}$ and $\Omega \in U_{\alpha}(\mathbf{R}^n)$, we have $d_{\alpha,\Omega}(x, y) \leq C_{\alpha,\Omega}\|x - y\|^{\alpha}$ so that

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) < 2C_{\alpha,\Omega}\|x - y\|^{\alpha}.$$

By Lemma 2.1, part (i),

$$\text{length}(\gamma) \leq 2e^{2C_{\alpha,\Omega}}\|x - y\|$$

proving (i). Hence,

$$\text{length}(\gamma) \leq 2e^{2C_{\alpha,\Omega}}\theta = 2e^{2C_{\alpha,\Omega}}(\frac{1}{2}e^{-2C_{\alpha,\Omega}})\theta_{\alpha,\Omega} = \theta_{\alpha,\Omega}$$

so that for every $u, v \in \gamma$ we have $\|u - v\| \leq \text{length}(\gamma) \leq \theta_{\alpha,\Omega}$. Since $\Omega \in U_{\alpha}(\mathbf{R}^n)$, this implies

$$d_{\alpha,\Omega}(u, v) \leq C_{\alpha,\Omega}\|u - v\|^{\alpha} \leq C_{\alpha,\Omega} \text{length}^{\alpha}(\gamma_{uv}).$$

Combining this inequality with (2.14) and (2.13), we obtain

$$\begin{aligned} \int_{\gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) &\leq C_{\alpha,\Omega} \text{length}^{\alpha}(\gamma_{uv}) + \delta \\ &\leq C_{\alpha,\Omega} \text{length}^{\alpha}(\gamma_{uv}) + \text{length}^{\alpha}(\gamma_{uv}) \end{aligned}$$

proving (ii) and the lemma. $\square$

Put

$$m := [2(2C_{\alpha,\Omega})^{\frac{1}{1-\alpha}}] + 1. \quad (2.15)$$

Let k be a positive integer such that

$$2e^{2C_{\alpha,\Omega}}\|x - y\|(1 - 1/m)^k \leq \varepsilon. \quad (2.16)$$

Finally, we put

$$\delta := \min\{d_{\alpha,\Omega}(x, y), m^{-\alpha k}\|x - y\|^{\alpha}\}.$$

Thus $0 < \delta \leq d_{\alpha,\Omega}(x, y)$ and

$$\|x - y\|m^{-k} \geq \delta^{\frac{1}{\alpha}}. \quad (2.17)$$

Let $\Gamma \subset \Omega$ be a rectifiable curve joining x to y such that

$$\int_{\Gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) < d_{\alpha,\Omega}(x, y) + \delta.$$

Then, by Lemma 2.7,

$$\int_{\Gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq 2C_{\alpha,\Omega} \|x - y\|^{\alpha} \quad (2.18)$$

and

$$\text{length}(\Gamma) \leq 2e^{2C_{\alpha,\Omega}} \|x - y\|. \quad (2.19)$$

Moreover, for every $u, v \in \Gamma$ such that

$$\text{length}(\Gamma_{uv}) \geq \delta^{\frac{1}{\alpha}} \quad (2.20)$$

the following inequality

$$\int_{\Gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq 2C_{\alpha,\Omega} \text{length}^{\alpha}(\Gamma_{uv}) \quad (2.21)$$

holds.

Inequality (2.21) and part (ii) of Lemma 2.2 imply:

$$\frac{1}{\text{length}(\Gamma_{uv})} \int_{\Gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq 4C_{\alpha,\Omega} \inf_{z \in \Gamma_{uv}} \text{dist}(z, \partial\Omega)^{\alpha-1}. \quad (2.22)$$

Put

$$L := \text{length}(\Gamma).$$

Since $\|x - y\| \leq L$, by (2.17) and (2.20), for every $u, v \in \Gamma$ such that

$$\text{length}(\Gamma_{uv}) \geq L m^{-k} \quad (2.23)$$

inequality (2.21) is satisfied.

By $\mathcal{I}_m$ we denote the family of all m -adic closed subintervals of the interval $I_0 := [0, L]$. recall that this family of intervals can be obtained by the standard iterative procedure: we start with the entire interval $[0, L]$ and, at each level of the construction, we split every interval of the given level into m equally sized closed subintervals.

Let

$$\mathcal{I}_{j,m} := \{\text{all } m\text{-adic intervals of the } j\text{-th level}\}.$$

Thus $\mathcal{I}_{0,m} := \{[0, L]\}$, $\mathcal{I}_{1,m} := \{[Li/m, L(i+1)/m] : i = 0, 1, \dots, m-1\}$ etc. Clearly, $|I| = L m^{-j}$ for every interval I of the j -th level. Put

$$S_{k,m} := \bigcup_{j=0}^k \mathcal{I}_{j,m} = \{\text{all } m\text{-adic intervals of the level at most } k\}.$$

Let us parameterize Γ by arclength; thus we identify γ with a function $\Gamma : [0, L] \rightarrow \Omega$ satisfying $\Gamma(0) = x, \Gamma(L) = y$. Finally, put

$$g(t) := \text{dist}(\Gamma(t), \partial\Omega)^{\alpha-1}, \quad t \in [0, L].$$

Then, by (2.23) and (2.22), the following is true: For each m -adic interval $I \in S_{k,m}$ we have

$$\frac{1}{|I|} \int_I g(t) dt \leq C_g \inf_{t \in I} g(t) \quad (2.24)$$

with $C_g := 4C_{\alpha,\Omega}$.

Following Melas [Mel] we say that g is a Muckenhoupt A_1 -weight on $[0, L]$ with respect to the family $S := S_{k,m}$ of all m -adic intervals of the level at most k . We let $\mathcal{M}_S$ denote the corresponding maximal operator for the family S :

$$\mathcal{M}_S g(t) := \sup \left\{ \frac{1}{|I|} \int_I |g(t)| dt : I \ni t, I \in S \right\}. \quad (2.25)$$

Thus (2.24) is equivalent to the inequality

$$\mathcal{M}_S g(t) \leq C_g g(t), \quad t \in [0, L].$$

Put

$$q^\sharp := \frac{\log m}{\log(m - (m-1)/C_g)}$$

and $q^* := (1 + q^\sharp)/2$. Clearly, $1 \leq C_g < \infty$ so that $q^\sharp, q^* > 1$.

We will be needed the following corollary of a general result proved in [Mel].

Theorem 2.8 *For any A_1 -weight g (with respect to S) and any q , $1 \leq q \leq q^*$, the following inequality*

$$\left(\frac{1}{L} \int_0^L (\mathcal{M}_S g)^q dt \right)^{\frac{1}{q}} \leq \tilde{C} \left(\frac{1}{L} \int_0^L g dt \right) \quad (2.26)$$

holds. Here $\tilde{C}$ is a constant depending only on m and C_g .

Remark 2.9 Actually the theorem is true for $1 \leq q < q^\sharp$ but with $\tilde{C}$ depending on m, C_g and q , see [Mel].

Corollary 2.10 *For any A_1 -weight g (with respect to S), any family $\mathcal{A}$ of non-overlapping m -adic intervals of the level at most k and any q , $1 \leq q \leq q^*$, we have*

$$\left(\frac{1}{L} \sum_{I \in \mathcal{A}} \left(\frac{1}{|I|} \int_I g dt \right)^q |I| \right)^{\frac{1}{q}} \leq \tilde{C} \left(\frac{1}{L} \int_0^L g dt \right). \quad (2.27)$$

Proof. In fact, by definition (2.25), for every $I \in \mathcal{A}$ and every $t \in I$

$$\frac{1}{|I|} \int_I g \, ds \leq \mathcal{M}_S g(t)$$

so that

$$\left(\frac{1}{|I|} \int_I g \, dt \right)^q |I| \leq \int_I (\mathcal{M}_S g)^q \, dt.$$

Therefore the left-hand side of (2.27) does not exceed

$$\left(\frac{1}{L} \sum_{I \in \mathcal{A}} \int_I (\mathcal{M}_S g)^q \, dt \right)^{\frac{1}{q}} \leq \left(\frac{1}{L} \int_0^L (\mathcal{M}_S g)^q \, dt \right)^{\frac{1}{q}}$$

which together with (2.26) implies the required inequality (2.27). $\square$

We turn to construction of a family $\mathcal{A} \subset S_{k,m}$ of non-overlapping m -adic intervals of the level at most k such that for each $I \in \mathcal{A}$

$$\sup_I g \leq C \inf_I g$$

and

$$|[0, L] \setminus \{\cup I : I \in \mathcal{A}\}| < \varepsilon.$$

Here C is a constant depending only on n, α , and $C_{\alpha, \Omega}$.

Let $I = [t_1, t_2] \in S_{k-1,m}$ be an m -adic interval of the level at most $k-1$ and let $u := \Gamma(t_1), v := \Gamma(t_2)$. By (2.21) and part (i) of Lemma 2.2, there exists $t_I \in I$ such the point $z_I = \Gamma(t_I) \in \Gamma_{uv}$ satisfies the following inequality:

$$\text{length}(\Gamma_{uv}) \leq C' \text{dist}(z_I, \partial\Omega) \quad (2.28)$$

with $C' := (2C_{\alpha, \Omega})^{\frac{1}{1-\alpha}}$.

Let us split the interval I into m equal subintervals $I^{(1)}, \dots, I^{(m)}$. Then $t_I \in I^{(j)}$ for some $j \in \{1, \dots, m\}$. By $\Gamma^{(j)} := \Gamma|_{I^{(j)}}$ we denote the arc corresponding to the interval $I^{(j)}$. Thus $I^{(j)} \in S_{k,m}$ is an m -adic interval of the level at most k and

$$\text{length}(\Gamma^{(j)}) = |I^{(j)}| = |I|/m = \text{length}(\Gamma_{uv})/m.$$

Since $\text{dist}(\cdot, \partial\Omega)$ is a Lipschitz function, for every $t \in I^{(j)}$ we have

$$|\text{dist}(z_I, \partial\Omega) - \text{dist}(z(t), \partial\Omega)| \leq \|z_I - z(t)\| \leq \text{length}(\Gamma^{(j)}) = \text{length}(\Gamma_{uv})/m.$$

Combining this inequality with (2.28) we obtain

$$|\text{dist}(z_I, \partial\Omega) - \text{dist}(z(t), \partial\Omega)| \leq \frac{C'}{m} \text{dist}(z_I, \partial\Omega).$$

But $m := [2C'] + 1$, see (2.15), so that

$$|\text{dist}(z_I, \partial\Omega) - \text{dist}(z(t), \partial\Omega)| \leq \frac{1}{2} \text{dist}(z_I, \partial\Omega).$$

Hence

$$\frac{1}{2} \text{dist}(z_I, \partial\Omega) \leq \text{dist}(z(t), \partial\Omega) \leq \frac{3}{2} \text{dist}(z_I, \partial\Omega), \quad t \in I^{(j)}.$$

We let $\tilde{I}$ denote the interval $I^{(j)}$. Thus we have proved that for each $I \in S_{k-1,m}$ there exists a subinterval $\tilde{I} \in S_{k,m}$, $\tilde{I} \subset I$, such that

$$\max_{t \in \tilde{I}} \text{dist}(z(t), \partial\Omega) \leq 3 \min_{t \in \tilde{I}} \text{dist}(z(t), \partial\Omega).$$

Since $g(t) := \text{dist}(z(t), \partial\Omega)^{\alpha-1}$, we obtain

$$\max_{t \in \tilde{I}} g(t) \leq 3^{1-\alpha} \min_{t \in \tilde{I}} g(t).$$

Now we construct the family $\mathcal{A} \subset S_{k,m}$ as follows. At the first stage for the interval $I_0 := [0, L]$ we determine an m -adic interval $\tilde{I}_0 \in \mathcal{I}_{1,m}$ of the first level and put $\mathcal{A}_1 := \{\tilde{I}_0\}$ and $U_1 := \tilde{I}_0$.

Let us consider the set $[0, L] \setminus U_1$ which consists of $m-1$ m -adic intervals of the first level. We let $\mathcal{B}_1$ denote the family of these intervals. For every $I \in \mathcal{B}_1$ we construct the interval $\tilde{I} \in \mathcal{I}_{2,m}$ and put

$$\mathcal{A}_2 := \{\tilde{I} \in \mathcal{I}_{2,m} : I \in \mathcal{B}_1\}.$$

By U_2 we denote the set

$$U_2 := U_1 \cup \{I : I \in \mathcal{A}_2\}.$$

Now the set $[0, L] \setminus U_2$ consists of $(m-1)^2$ m -adic intervals of the second level. We denote the family of these intervals by $\mathcal{B}_2$ and finish the second stage of the procedure.

After the k -th stages of this procedure we obtain the families $\mathcal{A}_j \subset \mathcal{I}_{j,m}$, $j = 1, 2, \dots, k$, of m -adic intervals. We put

$$\mathcal{A} = \cup \{\mathcal{A}_j : j = 1, \dots, k\}.$$

Thus $\mathcal{A} \subset S_{k,m}$ is a family of m -adic intervals of the level at most k . We know that for every interval $I \in \mathcal{A}$ the following inequality

$$\max_{t \in I} g(t) \leq 3^{1-\alpha} \min_{t \in I} g(t) \tag{2.29}$$

holds. We also know that the set

$$U = U_k := \cup \{I : I \in \mathcal{A}\}$$

has the following property: the set

$$E := [0, L] \setminus U = [0, L] \setminus \cup \{I : I \in \mathcal{A}\}$$

consists of $(m-1)^k$ m -adic intervals of the k -th level. Since $|I| = m^{-k} L$ for each $I \in \mathcal{I}_{k,m}$, we obtain

$$|E| = \frac{(m-1)^k}{m^k} L.$$

But, by (2.19),

$$L = \text{length}(\Gamma) \leq C'' \|x - y\|$$

where $C'' := 2e^{2C_{\alpha,\Omega}}$. Hence,

$$|E| \leq C'' \|x - y\| (1 - 1/m)^k.$$

Combining this inequality with (2.16), we obtain the required estimate

$$|E| = |[0, L] \setminus U| \leq \varepsilon. \quad (2.30)$$

Now for the family $\mathcal{A}$ constructed above let us estimate from below the quantity

$$T := \left(\frac{1}{L} \sum_{I \in \mathcal{A}} \left(\frac{1}{|I|} \int_I g \, dt \right)^q |I| \right)^{\frac{1}{q}}$$

from the left-hand side of inequality (2.27). By (2.29), for each $I \in \mathcal{A}$ we have

$$\int_I g^q \, dt \leq |I| \max_{t \in I} g^q(t) \leq 3^{q(1-\alpha)} |I| (\min_{t \in I} g(t))^q \leq 3^{q(1-\alpha)} |I| \left(\frac{1}{|I|} \int_I g \, dt \right)^q$$

so that

$$T^q \geq 3^{q(\alpha-1)} \frac{1}{L} \sum_{I \in \mathcal{A}} \int_I g^q \, dt = 3^{q(\alpha-1)} \frac{1}{L} \int_U g^q \, dt.$$

(Recall that $U = \cup \{I : I \in \mathcal{A}\}$.)

By Corollary 2.10,

$$T \leq \tilde{C} \left(\frac{1}{L} \int_0^L g \, dt \right)$$

so that

$$\frac{1}{L} \int_U g^q \, dt \leq 3^{q(1-\alpha)} T^q \leq C_1 \left(\frac{1}{L} \int_0^L g \, dt \right)^q.$$

with $C_1 := 3^{q(\alpha-1)} \tilde{C}$. On the other hand, by inequality (2.18),

$$\int_0^L g \, dt = \int_{\Gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} \, ds(z) \leq 2C_{\alpha,\Omega} \|x - y\|^\alpha.$$

Hence

$$\frac{1}{L} \int_U g^q dt \leq C_1 \left(\frac{1}{L} \int_0^L g dt \right)^q \leq C_2 \|x - y\|^{q\alpha} / L^q$$

with $C_2 := (2C_{\alpha,\Omega})^q C_1$.

Recall that this inequality holds for all $q \in [1, q^*]$, see Corollary 2.10. We put $\tilde{q} := \min\{q^*, \frac{1-\alpha/2}{1-\alpha}\}$ and $\alpha^* := 1 - \tilde{q}(1 - \alpha)$. Since $q^* > 1$ and $0 < \alpha < 1$, we have $1 < \tilde{q} \leq q$ and $0 < \alpha^* < \alpha$.

Let $\tau \in [\alpha^*, \alpha]$ and let $q := \frac{1-\tau}{1-\alpha}$. Then $q \in [1, q^*]$ so that

$$\int_U \text{dist}(\Gamma(t), \partial\Omega)^{\tau-1} dt = \int_U g^{\frac{\tau-1}{\alpha-1}}(t) dt = \int_U g^q(t) dt \leq \int_0^L g^q(t) dt \leq C_2 \|x - y\|^{q\alpha} / L^{q-1}$$

Since $\|x - y\| \leq L$, we obtain

$$\int_U \text{dist}(\Gamma(t), \partial\Omega)^{\tau-1} dt \leq C_2 \|x - y\|^{q\alpha} / L^{q-1} \leq C_2 \|x - y\|^{q\alpha - q + 1} = C_2 \|x - y\|^\tau. \quad (2.31)$$

Finally, we put

$$\tilde{\Gamma} := \Gamma|_U.$$

Then inequalities (2.18), (2.30) and (2.31) shows that inequalities (1.10), (1.9) and (1.7) of Theorem 1.4 are satisfied.

It remains to prove inequality (1.8). We observe that the set $U = U_k$ is obtained by the standard Cantor procedure (for m -adic intervals; recall that in the classical case $m = 3$). The reader can easily see that this Cantor set possess the following property: for each interval I centered in U with $|I| \leq 2L$ we have

$$|I| \leq 4m |I \cap U|. \quad (2.32)$$

Let $B = B(c, r)$ be a ball of radius r centered at a point $c \in \tilde{\Gamma}$. We may assume that $r \leq \|x - y\|/4$. Then either x or y does not belong to B . Suppose that $y \notin B$. Then there exists a point $v \in \partial B \cap \Gamma_{cy}$ such that the arc $\Gamma_{cv} \subset B$.

Let us consider the arc Γ_{xc} . If $\Gamma_{xc} \subset B$, we put

$$\ell := \text{length}(\Gamma_{cv}). \quad (2.33)$$

Clearly, in this case $\ell \geq r$.

Assume that $\Gamma_{xc} \not\subset B$. Then there exists a point $u \in \partial B \cap \Gamma_{xc}$ such that the arc $\Gamma_{uc} \subset B$. In this case we put

$$\ell := \min\{\text{length}(\Gamma_{uc}), \text{length}(\Gamma_{cv})\}. \quad (2.34)$$

Since $\text{length}(\Gamma_{uc}) \geq r$, $\text{length}(\Gamma_{cv}) \geq r$, again we have $\ell \geq r$.

Recall that $c \in \tilde{\Gamma}$ so that $c = \Gamma(A)$ for some $a \in U$. By I we denote the interval $I := [a - \ell, a + \ell]$. Then, by definitions (2.33), (2.34), the arc $\Gamma(I) \subset B$. Since I is centered in U and $|I| = 2\ell \leq 2L$, by (2.32),

$$|I| \leq 4m |I \cap U|.$$

Since $\Gamma(I) \subset B$, we have $|I \cap U| \leq \text{length}(B \cap \tilde{\Gamma})$ so that

$$\text{diam } B = 2r \leq 2\ell = |I| \leq 4m |I \cap U| \leq 4m \text{length}(B \cap \tilde{\Gamma})$$

proving (1.8).

Theorem 1.4 is completely proved. $\square$

3. Sobolev functions on subhyperbolic domains.

Let us fix some additional notation. In what follows, the terminology “cube” will mean a closed cube in $\mathbf{R}^n$ whose sides are parallel to the coordinate axes. We let $Q(x, r)$ denote the cube in $\mathbf{R}^n$ centered at x with side length $2r$. Given $\lambda > 0$ and a cube Q we let λQ denote the dilation of Q with respect to its center by a factor of λ . (Thus $\lambda Q(x, r) = Q(x, \lambda r)$.)

It will be convenient for us to measure distances in $\mathbf{R}^n$ in the uniform norm

$$\|x\| := \max\{|x_i| : i = 1, \dots, n\}, \quad x = (x_1, \dots, x_n) \in \mathbf{R}^n.$$

Thus every cube

$$Q = Q(x, r) := \{y \in \mathbf{R}^n : \|y - x\| \leq r\}$$

is a “ball” in $\|\cdot\|$ -norm of “radius” r centered at x .

By χ_A we denote the characteristic function of A ; we put $\chi_A \equiv 0$ whenever $A = \emptyset$.

Let $\mathcal{A} = \{Q\}$ be a family of cubes in $\mathbf{R}^n$. By $M(\mathcal{A})$ we denote its *covering multiplicity*, i.e., the minimal positive integer M such that every point $x \in \mathbf{R}^n$ is covered by at most M cubes from $\mathcal{A}$. Thus

$$M(\mathcal{A}) := \sup_{x \in \mathbf{R}^n} \sum_{Q \in \mathcal{A}} \chi_Q(x).$$

Recall that $\mathcal{P}_k$ denotes the space of polynomials of degree at most k defined on $\mathbf{R}^n$. Also recall that, given a k -times differentiable function F and a point $x \in \mathbf{R}^n$, we let $T_x^k(F)$ denote the Taylor polynomial of F at x of degree at most k :

$$T_x^k(F)(y) := \sum_{|\beta| \leq k} \frac{1}{\beta!} (D^\beta F)(x) (y - x)^\beta, \quad y \in \mathbf{R}^n.$$

Let Ω be a domain in $\mathbf{R}^n$ and let $k \in \mathbf{N}$ and $p \in [1, \infty]$. We let $L_p^k(\Omega)$ denote the (homogeneous) Sobolev space of all functions $f \in L_{1,loc}(\Omega)$ whose distributional partial derivatives on Ω of order k belong to $L_p(\Omega)$. $L_p^k(\Omega)$ is normed by

$$\|f\|_{L_p^k(\Omega)} := \left(\int_{\Omega} \|\nabla^k f\|^p dx \right)^{\frac{1}{p}}$$

where $\nabla^k f$ denotes the vector with components $D^\beta f$, $|\beta| = k$, and

$$\|\nabla^k f\|(x) := \left(\sum_{|\beta|=k} |D^\beta f(x)|^2 \right)^{\frac{1}{2}}, \quad x \in \Omega.$$

By the Sobolev imbedding theorem, see e.g., [M], p. 60, every $f \in L_p^k(\Omega)$, $p > n$, can be redefined, if necessary, in a set of Lebesgue measure zero so that it belongs to the space $C^{k-1}(\Omega)$. Moreover, for every cube $Q \subset \Omega$, every $x, y \in Q$ and every multiindex β , $|\beta| \leq k-1$, the following inequality

$$|D^\beta T_x^{k-1}(f)(x) - D^\beta T_y^{k-1}(f)(x)| \leq C(n, p) \|x - y\|^{k-|\beta|-\frac{n}{p}} \left(\int_Q \|\nabla^k f\|^p dx \right)^{\frac{1}{p}} \quad (3.1)$$

holds. In particular, the partial derivatives of order $k-1$ satisfy a (local) Hölder condition of order $\alpha := 1 - \frac{n}{p}$:

$$|D^\beta f(x) - D^\beta f(y)| \leq C(n, p) \|f\|_{L_p^k(\Omega)} \|x - y\|^{1-\frac{n}{p}}, \quad |\beta| = k-1,$$

provided Q is a cube in Ω and $x, y \in Q$.

Thus, for $p > n$, we can identify each element $f \in L_p^k(\Omega)$ with its unique C^{k-1} -representative on Ω . This will allow us to restrict our attention to the case of Sobolev C^{k-1} -functions.

The main result of this section is the following

Theorem 3.1 *Let $n < p < \infty$, $\alpha = (p-n)/(p-1)$, and let Ω be an α -subhyperbolic domain in $\mathbf{R}^n$. There exists a constant $p^*, n < p^* < p$, and constants $\lambda, \theta, C > 0$ depending only on $n, p, k, C_{\alpha, \Omega}$ and $\theta_{\alpha, \Omega}$, such that the following is true: Let $f \in L_p^k(\Omega)$, $x, y \in \Omega$, $\|x - y\| \leq \theta$, and let $Q_{xy} := Q(x, \|x - y\|)$. Then for every multiindex β , $|\beta| \leq k-1$, the following inequality*

$$|D^\beta T_x^{k-1}(f)(x) - D^\beta T_y^{k-1}(f)(x)| \leq C \|x - y\|^{k-|\beta|-\frac{n}{p^*}} \left(\int_{(\lambda Q_{xy}) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \quad (3.2)$$

holds.

Proof. We will be needed the following

Lemma 3.2 *Let $x, y \in \mathbf{R}^n$ and let $\gamma \subset \Omega$ be a continuous curve joining x to y . There is a finite family of cubes $\mathcal{Q} = \{Q_0, \dots, Q_m\}$ such that:*

(i). $Q_0 \ni x, Q_m \ni y$, $Q_i \neq Q_j$, $i \neq j$, $0 \leq i, j \leq m$, and

$$Q_i \cap Q_{i+1} \neq \emptyset, \quad i = 0, \dots, m-1.$$

(ii). For every cube $Q = Q(z, r) \in \mathcal{Q}$ we have $z \in \gamma$ and $r = \frac{1}{8} \text{dist}(z, \partial\Omega)$.

(iii). For each $Q \in \mathcal{Q}$ the cube $2Q \subset \Omega$. Moreover, the covering multiplicity of the family of cubes $2\mathcal{Q} := \{2Q : Q \in \mathcal{Q}\}$ is bounded by a constant $C = C(n)$.

Proof. For every $z \in \Gamma$ we let $Q^{(z)}$ denote the cube

$$Q^{(z)} := Q(z, \frac{1}{8} \text{dist}(z, \partial\Omega)).$$

We put $\mathcal{A} := \{Q^{(z)} : z \in \Gamma\}$. By the Besicovitch covering theorem, see e.g. [G], there exists a finite subcollection $\mathcal{B} \subset \mathcal{A}$ such that $\mathcal{B}$ still covers Γ but no point which lies in more than $C(n)$ of the cubes of $\mathcal{B}$. (Thus the covering multiplicity $M(\mathcal{B}) \leq C(n)$.)

Given $Q', Q'' \in \mathcal{B}$ we write $Q' \sim Q''$ if there exists a family $\{K_0, \dots, K_\ell\} \subset \mathcal{B}$ such that $K_0 = Q'$, $K_\ell = Q''$, $K_i \neq K_j, i \neq j$, $0 \leq i, j \leq \ell$, and $K_i \cap K_{i+1} \neq \emptyset$ for every $i = 0, \dots, \ell - 1$. Fix a cube $\tilde{Q} \in \mathcal{B}$ such that $x \in \tilde{Q}$, and put

$$\mathcal{B}' := \{Q \in \mathcal{B} : Q \sim \tilde{Q}\} \quad (3.3)$$

and $E := \cup\{Q : Q \in \mathcal{B}'\}$.

Prove that $y \in E$. Assume that this is not true, i.e., $y \notin E$. Since $\tilde{Q} \ni x$, we have $x \in E$ so that $E \cap \gamma \neq \emptyset$. Put $F := \cup\{Q : Q \in \mathcal{B} \setminus \mathcal{B}'\}$. Observe that every cube $Q \in \mathcal{B}$ such that $Q \cap E \neq \emptyset$ belongs to $\mathcal{B}'$ so that $E \cap F = \emptyset$.

On the other hand, since $\gamma \subset E \cup F$ and $y \notin E$, we have $y \in F$ proving that $F \cap \gamma \neq \emptyset$. Thus the sets $E \cap \gamma$ and $F \cap \gamma$ is a partition of the continuous curve γ into two closed disjoint sets; a contradiction. We have proved that $y \in Q^*$ for some $Q^* \in \mathcal{B}'$ so that, by definition (3.3), there exists a family of cubes $\mathcal{Q} = \{Q_0, \dots, Q_m\}$ satisfying conditions (i) and (ii) of the lemma.

Prove (iii). Let $Q = Q(z, r) \in \mathcal{Q}$. Then $z \in \gamma$ and $r = \frac{1}{8} \text{dist}(z, \partial\Omega)$ so that for every $u \in 2Q = Q(z, 2r)$ we have

$$\text{dist}(z, \partial\Omega) \leq \text{dist}(u, \partial\Omega) + \|u - z\| \leq \text{dist}(u, \partial\Omega) + 2r \leq \text{dist}(u, \partial\Omega) + \frac{1}{4} \text{dist}(z, \partial\Omega).$$

Hence,

$$0 < \frac{3}{4} \text{dist}(z, \partial\Omega) \leq \text{dist}(u, \partial\Omega)$$

proving that $2Q \subset \Omega$.

It remains to prove that the covering multiplicity $M(2\mathcal{Q}) \leq C(n)$. We know that $M(\mathcal{Q}) \leq M(\mathcal{B}) \leq C(n)$. Fix a cube $Q = Q(z, r) \in \mathcal{Q}$. Let $Q_i = Q(z_i, r_i) \in \mathcal{Q}$ be an arbitrary cube such that

$$(2Q) \cap (2Q_i) \neq \emptyset. \quad (3.4)$$

Then $\|z - z_i\| \leq 2r + 2r_i$ so that

$$r = \frac{1}{8} \text{dist}(z, \partial\Omega) \leq \frac{1}{8} \text{dist}(z_i, \partial\Omega) + \frac{1}{8} \|z - z_i\| \leq r_i + \frac{1}{8} (2r + 2r_i) = \frac{1}{4} r + \frac{5}{4} r_i.$$

Hence $r \leq \frac{5}{3} r_i$. In the same way we prove that $r_i \leq \frac{5}{3} r$.

Since $\mathcal{Q}$ has the covering multiplicity at most $C(n)$, this collection of cubes can be partitioned into at most $N(n)$ families of pairwise disjoint cubes, see e.g. [BrK]. Therefore, without loss of generality, we may assume that $\mathcal{Q}$ itself is a collection of pairwise disjoint cubes.

Since $\frac{1}{2}r \leq r_i \leq 2r$, we have $2^{-n}|Q| \leq |Q_i| \leq 2^n|Q|$. Also, by (3.4) and the inequality $r_i \leq 2r$, we have $Q_i \subset 7Q$. Thus the number of cubes Q_i satisfying (3.4) is bounded by $|7Q|/(2^{-n}|Q|) = 2^{n+3}$. The lemma is proved. $\square$

Let $x, y \in \Omega$, $\|x - y\| \leq \theta$, where θ is the constant from Theorem 1.4. By this theorem there exist constants $\alpha^* = \alpha^*(n, p)$, $0 < \alpha^* < \alpha$, and $C = C(n, p) > 0$ such that for every

$\varepsilon > 0$ there exists a rectifiable curve $\Gamma \subset \Omega$ and a finite family of arcs $\tilde{\Gamma} \subset \Gamma$ satisfying conditions (i),(ii) of the theorem.

Observe that, by inequality (1.7) (with $\tau = \alpha$) and by (1.10),

$$\int_{\Gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z) \leq C\|x - y\|^{\alpha}. \quad (3.5)$$

Also, by part (ii) of Theorem 1.4,

$$\text{length}(\Gamma) \leq \tilde{C}\|x - y\| \quad (3.6)$$

with $\tilde{C} = 2e^{2C_{\alpha,\Omega}}$, see (2.19).

By Lemma 3.2, there exists a collection of cubes

$$\mathcal{Q} = \{Q_0, \dots, Q_m\}$$

satisfying conditions (i)-(iii) of the lemma. In particular, by (i), $Q_0 \ni x, Q_m \ni y, Q_i \neq Q_j, i \neq j$, and

$$Q_i \cap Q_{i+1} \neq \emptyset, \quad i = 0, \dots, m-1.$$

Let $Q_i = Q(z_i, r_i), i = 0, \dots, m$, (recall that by (ii) $z_i \in \Gamma$ and $r_i = \frac{1}{8} \text{dist}(z_i, \partial\Omega)$). Let $a_i \in Q_{i-1} \cap Q_i, i = 1, \dots, m$. We also put $a_0 := x, a_{m+1} := y$.

We may assume that for every $Q \in \mathcal{Q}$ either $x \notin Q$ or $y \notin Q$. In fact, otherwise $x, y \in Q$. But by condition (iii) of Lemma 3.2, $2Q \subset \Omega$. Then the cube $Q(x, \|x - y\|) \subset 2Q \subset \Omega$ as well. It remains to apply inequality (3.1) to x and y with $p = p^*, n < p^* < p$, and (3.2) follows.

Thus we assume that for every cube $Q_i = Q(z_i, r_i) \in \mathcal{Q}$

$$\text{either } x \notin Q_i \text{ or } y \notin Q_i.$$

Since $x, y, z_i \in \Gamma$, we have $\partial Q_i \cap \Gamma \neq \emptyset$ so that there exists a point $a_i \in \partial Q_i \cap \Gamma$. Hence, by (3.6),

$$r_i = \|z_i - a_i\| \leq \text{length}(\Gamma \cap Q_i) \quad (3.7)$$

so that, by (3.6),

$$r_i \leq \text{length}(\Gamma) \leq \tilde{C}\|x - y\|, \quad i = 0, \dots, m. \quad (3.8)$$

Now we have

$$\begin{aligned} A &:= |D^{\beta}T_x^{k-1}(f)(x) - D^{\beta}T_y^{k-1}(f)(x)| = |D^{\beta}T_{a_0}^{k-1}(f)(a_0) - D^{\beta}T_{a_{m+1}}^{k-1}(f)(a_0)| \\ &\leq \sum_{i=0}^m |D^{\beta}T_{a_i}^{k-1}(f)(a_0) - D^{\beta}T_{a_{i+1}}^{k-1}(f)(a_0)|. \end{aligned}$$

Put

$$P_i(z) := T_{a_i}^{k-1}(f)(z) - T_{a_{i+1}}^{k-1}(f)(z), \quad i = 0, \dots, m.$$

The polynomial $P_i \in \mathcal{P}_{k-1}$ so that for every multiindex $\beta, |\beta| \leq k-1$, we have

$$D^\beta P_i(z) = \sum_{|\eta| \leq k-1-|\beta|} \frac{1}{\eta!} D^{\eta+\beta} P_i(a_i) (z - a_i)^\eta, \quad z \in \mathbf{R}^n.$$

Hence

$$|D^\beta P_i(a_0)| \leq C \sum_{|\eta| \leq k-1-|\beta|} |D^{\eta+\beta} P_i(a_i)| \|a_0 - a_i\|^{|\eta|}.$$

We put

$$\mathcal{Q}_1 := \{Q \in \mathcal{Q} : Q \cap \tilde{\Gamma} \neq \emptyset\}, \quad I_1 := \{i \in \{0, \dots, m\} : Q_i \in \mathcal{Q}_1\}, \quad (3.9)$$

and

$$\mathcal{Q}_2 := \mathcal{Q} \setminus \mathcal{Q}_1, \quad I_2 := \{i \in \{0, \dots, m\} : Q_i \in \mathcal{Q}_2\}. \quad (3.10)$$

Then

$$\begin{aligned} A &\leq \sum_{i=0}^m |D^\beta P_i(a_0)| \leq C \sum_{i=0}^m \sum_{|\eta| \leq k-1-|\beta|} |D^{\eta+\beta} P_i(a_i)| \|a_0 - a_i\|^{|\eta|} \\ &= C \sum_{|\eta| \leq k-1-|\beta|} \left(\sum_{i=0}^m |D^{\eta+\beta} P_i(a_i)| \|a_0 - a_i\|^{|\eta|} \right). \end{aligned}$$

Let

$$A'_\eta := \sum_{i \in I_1} |D^{\eta+\beta} P_i(a_i)| \|a_0 - a_i\|^{|\eta|}$$

and

$$A''_\eta := \sum_{i \in I_2} |D^{\eta+\beta} P_i(a_i)| \|a_0 - a_i\|^{|\eta|}.$$

We have proved that

$$A \leq C \sum_{|\eta| \leq k-1-|\beta|} (A'_\eta + A''_\eta). \quad (3.11)$$

Let $|\eta| \leq k-1-|\beta|$. Our next aim is to show that

$$A'_\eta \leq C \|x - y\|^{k-|\beta|-\frac{n}{p^*}} \left(\int_{(\lambda Q_{xy}) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \quad (3.12)$$

where

$$p^* := \frac{n - \alpha^*}{1 - \alpha^*}$$

and $\lambda := 2\tilde{C}$. (Recall that $Q_{xy} := Q(x, \|x - y\|)$.) Also we will prove that

$$A''_\eta \leq C \|x - y\|^{k-|\beta|-\frac{n}{p}} \left(\int_E \|\nabla^k f\|^p dx \right)^{\frac{1}{p}} \quad (3.13)$$

where E is a subset of Ω of the Lebesgue measure $|E| \leq C\varepsilon^n$.

Since

$$0 < \alpha^* = \frac{p^* - n}{p^* - 1} < \alpha = \frac{p - n}{p - 1} < 1,$$

we have $n < p^* < p$. Since $a_i, a_{i+1} \in Q_i = Q(z_i, r_i)$, $i \in I_1$, by inequality (3.1) (with $p = p^*$),

$$\begin{aligned} |D^{\eta+\beta} P_i(a_i)| &= |D^{\eta+\beta} T_{a_i}^{k-1}(f)(a_i) - D^{\eta+\beta} T_{a_{i+1}}^{k-1}(f)(a_i)| \\ &\leq C \|a_i - a_{i+1}\|^{k-|\eta|-|\beta|-\frac{n}{p^*}} \left(\int_{Q_i} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \end{aligned}$$

so that

$$|D^{\eta+\beta} P_i(a_i)| \leq C r_i^{k-|\eta|-|\beta|-\frac{n}{p^*}} \left(\int_{Q_i} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}, \quad i \in I_1. \quad (3.14)$$

In a similar way we prove that

$$|D^{\eta+\beta} P_i(a_i)| \leq C r_i^{k-|\eta|-|\beta|-\frac{n}{p}} \left(\int_{Q_i} \|\nabla^k f\|^p dx \right)^{\frac{1}{p}}, \quad i \in I_2.$$

Prove that for each $i \in \{0, \dots, m\}$ we have

$$\|a_0 - a_i\| \leq 2\tilde{C}\|x - y\|. \quad (3.15)$$

In fact,

$$\|a_0 - a_i\| = \|x - a_i\| \leq \|x - z_i\| + \|z_i - a_i\| \leq \|x - z_i\| + r_i.$$

Since $x, z_i \in \Gamma$, by (3.6), $\|x - z_i\| \leq \text{length}(\Gamma) \leq \tilde{C}\|x - y\|$. Also, by (3.8), $r_i \leq \tilde{C}\|x - y\|$, proving (3.15).

Hence, by (3.14),

$$A'_\eta \leq (2\tilde{C})^{|\eta|} \sum_{i \in I_1} |D^{\eta+\beta} P_i(a_i)| \|x - y\|^{|\eta|} \leq C \sum_{i \in I_1} r_i^{k-|\eta|-|\beta|-\frac{n}{p^*}} \|x - y\|^{|\eta|} \left(\int_{Q_i} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}.$$

Since $k - |\eta| - |\beta| \geq 1$ and $r_i \leq \tilde{C} \|x - y\|$, we have

$$\begin{aligned} r_i^{k-|\eta|-|\beta|-\frac{n}{p^*}} \|x - y\|^{|\eta|} &= r_i^{k-|\eta|-|\beta|-1} r_i^{1-\frac{n}{p^*}} \|x - y\|^{|\eta|} \\ &\leq C r_i^{1-\frac{n}{p^*}} \|x - y\|^{k-|\eta|-|\beta|-1} \|x - y\|^{|\eta|} \\ &= C r_i^{1-\frac{n}{p^*}} \|x - y\|^{k-|\beta|-1}. \end{aligned}$$

Hence

$$A'_\eta \leq C \|x - y\|^{k-|\beta|-1} \sum_{i \in I_1} r_i^{1-\frac{n}{p^*}} \left(\int_{Q_i} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}.$$

By the Hölder inequality,

$$A'_\eta \leq C \|x - y\|^{k-|\beta|-1} \left(\sum_{i \in I_1} (r_i^{1-\frac{n}{p^*}})^{\frac{p^*}{p^*-1}} \right)^{1-\frac{1}{p^*}} \left(\sum_{i \in I_1} \int_{Q_i} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}$$

so that

$$A'_\eta \leq C \|x - y\|^{k-|\beta|-1} \left(\sum_{i \in I_1} r_i^{\alpha^*} \right)^{1-\frac{1}{p^*}} \left(M(\mathcal{Q}_1) \int_{U_1} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}$$

where

$$U_1 := \cup \{Q : Q \in \mathcal{Q}_1\}.$$

Recall that $M(\mathcal{Q}_1)$ stands for the covering multiplicity of the collection $\mathcal{Q}_1$. Since $M(\mathcal{Q}_1) \leq M(\mathcal{Q}) \leq C(n)$, we obtain

$$A'_\eta \leq C \|x - y\|^{k-|\beta|-1} \left(\sum_{i \in I_1} r_i^{\alpha^*} \right)^{1-\frac{1}{p^*}} \left(\int_{U_1} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \quad (3.16)$$

In a similar way we prove that

$$A''_\eta \leq C \|x - y\|^{k-|\beta|-1} \left(\sum_{i \in I_2} r_i^\alpha \right)^{1-\frac{1}{p}} \left(\int_{U_2} \|\nabla^k f\|^p dx \right)^{\frac{1}{p}} \quad (3.17)$$

where

$$U_2 := \cup \{Q : Q \in \mathcal{Q}_2\}.$$

Let us prove that

$$\sum_{i \in I_1} r_i^{\alpha^*} \leq C \|x - y\|^{\alpha^*} \quad (3.18)$$

and

$$\sum_{i \in I_2} r_i^\alpha \leq C \|x - y\|^\alpha. \quad (3.19)$$

We begin with the proof of inequality (3.18). Let $i \in I_1$ and let

$$Q_i = Q(z_i, r_i), \quad r_i = \frac{1}{8} \text{dist}(z_i, \partial\Omega).$$

Recall that $Q_i \cap \tilde{\Gamma} \neq \emptyset$ so that there exist a point $b_i \in Q_i \cap \tilde{\Gamma}$. By (1.8),

$$r_i \leq C \text{length}(\tilde{\Gamma} \cap Q(b_i, r_i)).$$

But $Q(b_i, r_i) \subset 2Q_i = Q(z_i, 2r_i)$ so that

$$r_i \leq C \text{length}(2Q_i \cap \tilde{\Gamma}).$$

Hence,

$$r_i^{\alpha^*} \leq C r_i^{\alpha^*-1} \text{length}(2Q_i \cap \tilde{\Gamma}).$$

Let $z \in 2Q_i \cap \tilde{\Gamma}$. Then

$$|\text{dist}(z, \partial\Omega) - \text{dist}(z_i, \partial\Omega)| \leq \|z - z_i\| \leq 2r_i.$$

Since $\text{dist}(z_i, \partial\Omega) = 8r_i$, we have

$$\text{dist}(z, \partial\Omega) \leq \text{dist}(z_i, \partial\Omega) + 2r_i = 10r_i.$$

Hence,

$$r_i^{\alpha^*-1} \leq C \text{dist}(z, \partial\Omega)^{\alpha^*-1}, \quad z \in 2Q_i \cap \tilde{\Gamma},$$

so that

$$r_i^{\alpha^*} \leq C \int_{2Q_i \cap \tilde{\Gamma}} \text{dist}(z, \partial\Omega)^{\alpha^*-1} ds(z).$$

We put $2\mathcal{Q}_1 := \{2Q : Q \in \mathcal{Q}_1\}$. We have

$$\sum_{i \in I_1} r_i^{\alpha^*} \leq C \sum_{i \in I_1} \int_{2Q_i \cap \tilde{\Gamma}} \text{dist}(z, \partial\Omega)^{\alpha^*-1} ds(z)$$

so that

$$\sum_{i \in I_1} r_i^{\alpha^*} \leq C M(2\mathcal{Q}_1) \int_{\tilde{\Gamma}} \text{dist}(z, \partial\Omega)^{\alpha^*-1} ds(z).$$

By part (iii) of Lemma 3.2, the covering multiplicity

$$M(2\mathcal{Q}_1) \leq M(2\mathcal{Q}) \leq C(n).$$

Hence,

$$\sum_{i \in I_1} r_i^{\alpha^*} \leq C \int_{\tilde{\Gamma}} \text{dist}(z, \partial\Omega)^{\alpha^*-1} ds(z).$$

In a similar way we prove that

$$\sum_{i \in I_2} r_i^{\alpha} \leq C \int_{\Gamma} \text{dist}(z, \partial\Omega)^{\alpha-1} ds(z).$$

Combining these inequalities with (1.7) (where we put $\tau = \alpha^*$) and (3.5), we obtain the required inequalities (3.18) and (3.19).

Hence, by (3.16),

$$\begin{aligned} A'_\eta &\leq C \|x - y\|^{k-|\beta|-1} (\|x - y\|^{\alpha^*})^{1-\frac{1}{p^*}} \left(\int_{U_1} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \\ &= C \|x - y\|^{k-|\beta|-\frac{n}{p^*}} \left(\int_{U_1} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \end{aligned}$$

and, by (3.17),

$$A''_\eta \leq C \|x - y\|^{k-|\beta|-\frac{n}{p}} \left(\int_{U_2} \|\nabla^k f\|^p dx \right)^{\frac{1}{p}}. \quad (3.20)$$

Recall that for each $Q = Q(z, r) \in \mathcal{Q}$ its center, the point z , belongs to Γ . Moreover, by (3.8), $r \leq \tilde{C}\|x - y\|$, and, by (3.6), $\text{length}(\Gamma) \leq \tilde{C}\|x - y\|$. Hence,

$$Q \subset Q(x, 2\tilde{C}\|x - y\|) = \lambda Q_{xy}, \quad Q \in \mathcal{Q},$$

with $\lambda := 2\tilde{C}$. Now we have

$$U_1 := \cup \{Q : Q \in \mathcal{Q}_1\} \subset (\lambda Q_{xy}) \cap \Omega$$

so that

$$A'_\eta \leq C \|x - y\|^{k-|\beta|-\frac{n}{p^*}} \left(\int_{(\lambda Q_{xy}) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}$$

proving (3.12).

Let us put $E := U_2$ and prove that $|E| \leq C \varepsilon^n$. We have

$$|E| = |\cup \{Q : Q \in \mathcal{Q}_2\}| \leq \sum \{|Q| : Q \in \mathcal{Q}_2\} = 2^n \sum_{I \in I_2} r_i^n \leq 2^n \left(\sum_{I \in I_2} r_i \right)^n. \quad (3.21)$$

By (3.9) and (3.10),

$$Q_i \cap \tilde{\Gamma} = \emptyset \quad \text{for every } Q_i \in \mathcal{Q}_2,$$

so that

$$Q_i \cap \Gamma = Q_i \cap (\Gamma \setminus \tilde{\Gamma}).$$

But, by (3.7), $r_i \leq \text{length}(Q_i \cap \Gamma)$ so that $r_i \leq \text{length}(Q_i \cap (\Gamma \setminus \tilde{\Gamma}))$. Hence

$$\sum_{I \in I_2} r_i \leq \sum_{I \in I_2} \text{length}(Q_i \cap (\Gamma \setminus \tilde{\Gamma})) \leq M(\mathcal{Q}_2) \text{length}(\Gamma \setminus \tilde{\Gamma}).$$

Since $M(\mathcal{Q}_2) \leq M(\mathcal{Q}) \leq C(n)$, we obtain

$$\sum_{I \in I_2} r_i \leq C \text{length}(\Gamma \setminus \tilde{\Gamma})$$

so that, by (1.9),

$$\sum_{I \in I_2} r_i \leq C \varepsilon.$$

Combining this inequality with (3.21), we obtain the required inequality $|E| \leq C \varepsilon^n$. This inequality and (3.20) imply (3.13).

Now, by (3.11), (3.12) and (3.13),

$$\begin{aligned} A &:= |D^\beta T_x^{k-1}(f)(x) - D^\beta T_y^{k-1}(f)(x)| \leq C \sum_{|\eta| \leq k-1-|\beta|} (A'_\eta + A''_\eta) \\ &\leq C \sum_{|\eta| \leq k-1-|\beta|} \|x - y\|^{k-|\beta|-\frac{n}{p^*}} \left(\int_{(\lambda Q_{xy}) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \\ &\quad + C \sum_{|\eta| \leq k-1-|\beta|} \|x - y\|^{k-|\beta|-\frac{n}{p}} \left(\int_E \|\nabla^k f\|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

We obtain

$$\begin{aligned} A &\leq C \|x - y\|^{k-|\beta|-\frac{n}{p^*}} \left(\int_{(\lambda Q_{xy}) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \\ &\quad + C \|x - y\|^{k-|\beta|-\frac{n}{p}} \left(\int_E \|\nabla^k f\|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

But $f \in L_p^k(\Omega)$ so that

$$\int_{\Omega} \|\nabla^k f\|^p dx = \|f\|_{L_p^k(\Omega)}^p < \infty.$$

Hence,

$$\int_E \|\nabla^k f\|^p dx \rightarrow 0 \quad \text{as} \quad |E| = C\varepsilon^n \rightarrow 0,$$

proving the theorem. $\square$

4. Extension of Sobolev functions defined on subhyperbolic domains.

Given a cube $Q \subset \mathbf{R}^n$ and a function $f \in L_q(Q)$, $0 < q \leq \infty$, we let $\mathcal{E}_k(f; Q)_{L_q}$ denote the *normalized local best approximation* of f on Q in L_q -norm by polynomials of degree at most $k - 1$, see Brudnyi [Br1]. More explicitly, we define

$$\mathcal{E}_k(f; Q)_{L_q} := |Q|^{-\frac{1}{q}} \inf_{P \in \mathcal{P}_{k-1}} \|f - P\|_{L_q(Q)} = \inf_{P \in \mathcal{P}_{k-1}} \left(\frac{1}{|Q|} \int_Q |f - P|^q dx \right)^{\frac{1}{q}}.$$

In the literature $\mathcal{E}_k(f; Q)_{L_q}$ is also sometimes called the *local oscillation* of f , see e.g. Triebel [T2].

Given a locally integrable function f on $\mathbf{R}^n$, we define its *sharp maximal function* $f_k^\sharp$ by letting

$$f_k^\sharp(x) := \sup_{r>0} r^{-k} \mathcal{E}_k(f; Q(x, r))_{L_1}.$$

Recall that a function $f \in W_p^k(\mathbf{R}^n)$, $1 < p \leq \infty$, if and only if f and $f_k^\sharp$ are both in $L_p(\mathbf{R}^n)$, see Calderón [C]. Moreover, up to constants depending only on n, k and p the following equivalence,

$$\|f\|_{W_p^k(\mathbf{R}^n)} \sim \|f\|_{L_p(\mathbf{R}^n)} + \|f_k^\sharp\|_{L_p(\mathbf{R}^n)}, \quad (4.1)$$

holds.

This characterization motivates the following definition. Let S be a measurable subset of $\mathbf{R}^n$. Given a function $f \in L_{q,loc}(S)$, and a cube Q whose center is in S , we let $\mathcal{E}_k(f; Q)_{L_q(S)}$ denote the normalized best approximation of f on Q in $L_q(S)$ -norm:

$$\mathcal{E}_k(f; Q)_{L_q(S)} := |Q|^{-\frac{1}{q}} \inf_{P \in \mathcal{P}_k} \|f - P\|_{L_q(Q \cap S)} = \inf_{P \in \mathcal{P}_{k-1}} \left(\frac{1}{|Q|} \int_{Q \cap S} |f - P|^q dx \right)^{\frac{1}{q}}. \quad (4.2)$$

By $f_{k,S}^\sharp$, we denote the sharp maximal function of f on S ,

$$f_{k,S}^\sharp(x) := \sup_{r>0} r^{-k} \mathcal{E}_k(f; Q(x, r))_{L_1(S)}, \quad x \in S.$$

(Thus, $f_k^\sharp = f_{\alpha, \mathbf{R}^n}^\sharp$.)

Let $\Omega \subset \mathbf{R}^n$ be a subhyperbolic domain. The following two corollaries of Theorem 3.1 present estimates of the local best approximations and the sharp maximal function of a function $f \in W_p^k(\Omega)$ via the local L_p -norms and the maximal function of $\nabla^k f$.

Corollary 4.1 *Let $n < p < \infty$, $\alpha = (p - n)/(p - 1)$, and let Ω be an α -subhyperbolic domain in $\mathbf{R}^n$. There exists a constant $p^* \in (n, p)$ and constants $\theta, \lambda, C > 0$ depending only on $n, p, k, C_{\alpha, \Omega}$ and $\theta_{\alpha, \Omega}$, such that the following is true: Let $f \in L_p^k(\Omega)$. Then for every cube $Q = Q(x, r)$ with $x \in \Omega$ and $0 < r \leq \theta$ the following inequality*

$$r^{-k} \mathcal{E}_k(f; Q)_{L_\infty(\Omega)} \leq C \left(\frac{1}{|\lambda Q|} \int_{(\lambda Q) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}$$

holds.

Proof. Let p^* and θ be the constant from Theorem 3.1. Let $y \in Q(x, r)$ so that $\|y - x\| \leq r \leq \theta$. Applying Theorem 3.1 to the points y, x (with $\beta = 0$), we obtain

$$|f(y) - T_x^{k-1}(f)(y)| \leq C \|x - y\|^{k - \frac{n}{p^*}} \left(\int_{(\lambda Q_{xy}) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}.$$

Recall that $Q_{xy} := Q(x, \|x - y\|)$.

Since $n < p^*$, we have $k - \frac{n}{p^*} > 0$ so that

$$\begin{aligned} |f(y) - T_x^{k-1}(f)(y)| &\leq C r^{k - \frac{n}{p^*}} \left(\int_{(\lambda Q_{xy}) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \\ &\leq C r^k \left(\frac{1}{|\lambda Q|} \int_{(\lambda Q) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}}. \end{aligned}$$

Hence,

$$\begin{aligned} \mathcal{E}_k(f; Q)_{L_\infty(\Omega)} &:= \inf_{P \in \mathcal{P}_{k-1}} \sup_{y \in Q \cap \Omega} |f(y) - P(y)| \leq \sup_{y \in Q \cap \Omega} |f(y) - T_x^{k-1}(f)(y)| \\ &\leq C r^k \left(\frac{1}{|\lambda Q|} \int_{(\lambda Q) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \end{aligned}$$

proving the corollary. $\square$

Given a function g defined on Ω we let $g^\wedge$ denote its extension by zero to all of $\mathbf{R}^n$. Thus $g^\wedge(x) := g(x)$, $x \in \Omega$, and $g^\wedge(x) := 0$, $x \notin \Omega$.

As usual, given a function $u \in L_{1,loc}(\mathbf{R}^n)$ by $\mathcal{M}f$ we denote the Hardy-Littlewood maximal function

$$\mathcal{M}[f](x) := \sup_{t>0} \frac{1}{|Q(x, t)|} \int_{Q(x, t)} |f(y)| dy.$$

Corollary 4.2 *Let $n < p < \infty$, $\alpha = (p - n)/(p - 1)$, and let Ω be an α -subhyperbolic domain in $\mathbf{R}^n$. There exists a constant $p^*, n < p^* < p$, such that for every function $f \in L_p^k(\Omega)$ and every $x \in \Omega$ the following inequality*

$$f_{k,\Omega}^\sharp(x) \leq C \left\{ (\mathcal{M}[(\|\nabla^k f\|^\lambda)^{p^*}](x))^{\frac{1}{p^*}} + \mathcal{M}[f^\lambda](x) \right\}$$

holds. The constants p^* and C depend only on $n, p, k, C_{\alpha,\Omega}$ and $\theta_{\alpha,\Omega}$.

Proof. Let p^*, λ and θ be the constants from Corollary 4.1. By this corollary,

$$\begin{aligned} \sup_{0 < r \leq \theta} r^{-k} \mathcal{E}_k(f; Q(x, r))_{L_1(\Omega)} &\leq \sup_{0 < r \leq \theta} r^{-k} \mathcal{E}_k(f; Q(x, r))_{L_\infty(\Omega)} \\ &\leq C \sup_{0 < r \leq \theta} \left(\frac{1}{|Q(x, \lambda r)|} \int_{Q(x, \lambda r) \cap \Omega} \|\nabla^k f\|^{p^*} dx \right)^{\frac{1}{p^*}} \\ &\leq C \left\{ (\mathcal{M}[(\|\nabla^k f\|^\lambda)^{p^*}](x))^{\frac{1}{p^*}} \right\}. \end{aligned}$$

On the other hand, by (4.2),

$$\begin{aligned} \sup_{r > \theta} r^{-k} \mathcal{E}_k(f; Q(x, r))_{L_1(\Omega)} &\leq \theta^{-k} \sup_{r > \theta} \mathcal{E}_k(f; Q(x, r))_{L_1(\Omega)} \\ &\leq \theta^{-k} \sup_{r > \theta} \left(\frac{1}{|Q(x, r)|} \int_{Q(x, r) \cap \Omega} |f| dx \right) \\ &\leq \theta^{-k} \mathcal{M}(f^\lambda)(x) \end{aligned}$$

proving the lemma. $\square$

In [S4] we show that the restrictions of Sobolev functions to *regular subsets* of $\mathbf{R}^n$ can be described in a way similar to the Calderón's criterion (4.1), i.e., via L_p -norms of a function and its sharp maximal function on a set. We recall that a measurable set $S \subset \mathbf{R}^n$ is said to be *regular* if there are constants $\sigma_S \geq 1$ and $\delta_S > 0$ such that, for every cube Q with center in S and with diameter $\text{diam } Q \leq \delta_S$,

$$|Q| \leq \sigma_S |Q \cap S|.$$

Theorem 4.3 ([S4]) *Let S be a regular subset of $\mathbf{R}^n$. Then a function $f \in L_p(S)$, $1 < p \leq \infty$, can be extended to a function $F \in W_p^k(\mathbf{R}^n)$ if and only if its sharp maximal function $f_{k,S}^\sharp \in L_p(S)$. In addition,*

$$\|f\|_{W_p^k(\mathbf{R}^n)|_S} \sim \|f\|_{L_p(S)} + \|f_{k,S}^\sharp\|_{L_p(S)}$$

with constants of equivalence depending only on n, k, p, σ_S and δ_S .

Proof of Theorem 1.1. By inequality (1.1), Ω is an α -subhyperbolic domain with $\alpha = \frac{p-n}{p-1}$, so that, by Lemma 2.3, Ω is a *regular* subset of $\mathbf{R}^n$.

Let $p^* \in (n, p)$ be the constant from Corollary 4.2. Let $q > p^*$ and let $f \in W_q^k(\Omega)$. We have to prove that f can be extended to a function $F \in W_q^k(\mathbf{R}^n)$. Since Ω is regular, by Theorem 4.3 it suffices to show that the sharp maximal function $f_{k,\Omega}^\sharp$ belongs to $L_q(\Omega)$.

By Corollary 4.2,

$$f_{k,\Omega}^\sharp(x) \leq C \left\{ (\mathcal{M}[(\|\nabla^k f\|^\wedge)^{p^*}](x))^{\frac{1}{p^*}} + \mathcal{M}[f^\wedge](x) \right\}, \quad x \in \Omega,$$

so that

$$\begin{aligned} \|f_{k,\Omega}^\sharp\|_{L_q(\Omega)} &\leq C \left\{ \|(\mathcal{M}[(\|\nabla^k f\|^\wedge)^{p^*}])^{\frac{1}{p^*}}\|_{L_q(\Omega)} + \|\mathcal{M}[f^\wedge]\|_{L_q(\Omega)} \right\} \\ &\leq C \left\{ \|(\mathcal{M}[(\|\nabla^k f\|^\wedge)^{p^*}])^{\frac{1}{p^*}}\|_{L_q(\mathbf{R}^n)} + \|\mathcal{M}[f^\wedge]\|_{L_q(\mathbf{R}^n)} \right\}. \end{aligned}$$

By the Hardy-Littlewood maximal theorem

$$\|\mathcal{M}[f^\wedge]\|_{L_q(\mathbf{R}^n)} \leq C \|f^\wedge\|_{L_q(\mathbf{R}^n)} = C \|f\|_{L_q(\Omega)}.$$

(Recall that $f^\wedge$ denotes the extension of f by zero to all of $\mathbf{R}^n$.)

Applying the Hardy-Littlewood maximal theorem to the function $g := (\|\nabla^k f\|^\wedge)^{p^*}$ in the space $L_s(\mathbf{R}^n)$ with $s := q/p^* > 1$, we obtain

$$\begin{aligned} A := \|(\mathcal{M}[(\|\nabla^k f\|^\wedge)^{p^*}])^{\frac{1}{p^*}}\|_{L_q(\mathbf{R}^n)} &= (\|\mathcal{M}[g]\|_{L_s(\mathbf{R}^n)})^{\frac{1}{p^*}} \leq C (\|g\|_{L_s(\mathbf{R}^n)})^{\frac{1}{p^*}} \\ &= C \left\{ \left(\int_{\mathbf{R}^n} [(\|\nabla^k f\|^\wedge)^{p^*}]^{\frac{q}{p^*}} dx \right)^{\frac{p^*}{q}} \right\}^{\frac{1}{p^*}} \end{aligned}$$

so that

$$A \leq C \left(\int_{\mathbf{R}^n} (\|\nabla^k f\|^\wedge)^q dx \right)^{\frac{1}{q}} = C \left(\int_{\Omega} \|\nabla^k f\|^q dx \right)^{\frac{1}{q}} = C \|f\|_{L_q^k(\Omega)}.$$

Hence

$$\|f_{k,\Omega}^\sharp\|_{L_q(\Omega)} \leq C (\|f\|_{L_q^k(\Omega)} + \|f\|_{L_q(\Omega)}) \leq C \|f\|_{W_q^k(\Omega)}.$$

Theorem 1.1 is completely proved. $\square$

References

- [AHL] K. Astala, K. Hag, P. Hag, V. Lappalainen, Lipschitz Classes and the Hardy-Littlewood Property, *Mh. Math* 115 (1993), 267–279.
- [Br1] Yu. A. Brudnyi, Spaces that are definable by means of local approximations, *Trudy Moscov. Math. Obshch.*, 24 (1971) 69–132; English Transl.: *Trans. Moscow Math Soc.*, 24 (1974) 73–139.

- [BrK] Yu. A. Brudnyi, B.D. Kotljarskiĭ, A certain problem of combinatorial geometry, *Sibirsk. Mat. Z.* 11 (1970) 1171-1173; English transl. in *Siberian Math. J.* 11 (1970), 870-871.
- [BKos] S. Buckley, P. Koskela, Criteria for imbeddings of Sobolev-Poincaré type, *Internat. Math. Res. Notices* 18 (1996), 881–902.
- [BSt2] S. Buckley, A. Stanoyevitch, Weak slice conditions and Hölder imbeddings, *J. London Math. Soc.* 66 (2001), 690–706.
- [BSt3] S. Buckley, A. Stanoyevitch, Weak slice conditions, product domains, and quasiconformal mappings, *Rev. Math. Iberoam.* 17 (2001), 1–37.
- [BSt] S. Buckley, A. Stanoyevitch, Distinguishing properties of weak slice conditions, *Conformal geometry and dynamics*, *An Electronic Journal of the AMS* 7, (2003) 49-75.
- [BM] Yu. D. Burago, V. G. Maz'ya, Certain Questions of Potential Theory and Function Theory for Regions with Irregular Boundaries. (Russian) *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 3 1967, 152 pp.; English trans.: *Potential Theory and Function Theory for Irregular Regions. Seminars in Math., V. A. Steklov Math. Inst., Leningrad, Vol. 3, Consultants Bureau, New York* 1969 vii+68 pp.
- [C2] A. P. Calderón, Lebesgue spaces of differentiable functions and distributions, *Proc. Symp. Pure Math.*, vol. IV, (1961) pp. 33–49.
- [C] A. P. Calderón, Estimates for singular integral operators in terms of maximal functions, *Studia Math.* 44, (1972) 563–582 .
- [CF] R. Coifman and C. Fefferman, Weighted norm inequalities for maximal functions and singular integrals, *Studia Math.* 51 (1974) 241-250.
- [GR] J. Garcia-Cuerva, J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Mathematical Studies 116 (North-Holland, Amsterdam, 1985).
- [G] F. W. Gehring, The L^p integrability of the partial derivatives of a quasiconformal mapping, *Acta Math.* 130 (1973) 265-277.
- [GM] F.W. Gehring, O. Martio, Lipschitz classes and quasiconformal mappings, *Ann. Acad. Sci. Fenn. Ser. AI Math.* 10 (1985), 203-219.
- [GLV] V.M. Goldshtein, T.G. Latfullin, S.K. Vodopyanov, Criteria for extension of functions of the class L_2^1 from unbounded plain domains, *Siber. Math. J.*(Engl. transl.) 20, no. 2 (1979) 298-301.
- [GV1] V. M. Gol'dstein , S. K. Vodop'janov, Prolongement des fonctions de classe $\mathcal{L}_p^1$ et applications quasi conformes., *C. R. Acad. Sci. Paris Ser. A-B* 290 (1980), no. 10, A453–A456.

- [GV2] V. M. Gol'dstein and S. K. Vodop'janov, Prolongement des fonctions differentiables hors de domaines plans., C. R. Acad. Sci. Paris Ser. I Math. 293 (1981), no. 12, 581–584.
- [HKT] P. Hajlasz, P. Koskela, H. Tuominen, Sobolev embeddings, extensions and measure density condition, J. Funct. Anal. 254 (2008) 1217-1234.
- [HKT1] P. Hajlasz, P. Koskela, H. Tuominen, Measure density and extendability of Sobolev functions, Rev. Mat. Iberoamericana 24 (2008), no. 2, 645-669.
- [Jn] P. W. Jones, Quasiconformal mappings and extendability of functions in Sobolev spaces, Acta Math., 147 (1981), 71–78.
- [K] P. Koskela, Extensions and Imbeddings, J. Funct. Anal. 159 (1998) 369–383.
- [L] V. Lappalainen, Lip_h -extension domains, Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertations 56, (1985), 1-52.
- [Mac] B. Muckenhoupt, Weighted norm inequalities for the Hardy-Littlewood maximal function, Trans. Amer. Math. Soc. 165 (1972) 207-226.
- [M] V.G. Maz'ja, Sobolev spaces, Springer-Verlag, Berlin, 1985, xix+486 pp.
- [MP] V. Maz'ya, S. Poborchi, Differentiable Functions on Bad Domains, World Scientific, River Edge, NJ, 1997.
- [Mel] A. D. Melas, A sharp L_p inequality for dyadic A_1 weights in R^n , Bull. London Math. Soc., 37 (2005) 919-926.
- [S3] P. Shvartsman, On extension of Sobolev functions defined on regular subsets of metric measure spaces, J. Approx. Theory, 144 (2007), 139-161.
- [S4] P. Shvartsman, Local approximations and intrinsic characterizations of spaces of smooth functions on regular subsets of $\mathbf{R}^n$, Math. Nachr. 279 (2006), no.11, 1212-1241.
- [S7] P. Shvartsman, Sobolev W_p^1 -spaces on closed subsets of $\mathbf{R}^n$, Advances in Math. 220 (2009) 1842-1922.
- [St] E. M. Stein, Singular integrals and differentiability properties of functions, Princeton Univ. Press, Princeton, New Jersey, 1970.
- [T2] H. Triebel, Theory of function spaces. II. Monographs in Mathematics, 84. Birkhäuser Verlag, Basel, 1992.
- [W3] H. Whitney, Functions differentiable on the boundaries of regions, Ann. of Math. 35, no. 3, (1934), 482-485.
- [Zob1] N. Zobin, Whitney's problem on extendability of functions and an intrinsic metric, *Advances in Math.* **133** (1998) 96–132.

- [Zob2] N. Zobin, Extension of smooth functions from finitely connected planar domains, J. Geom. Anal. 9, no. 3, (1999), 489–509.

Department of Mathematics
Technion - Israel Institute of Technology
32000 Haifa
Israel

e-mail: pshv@tx.technion.ac.il